

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

AP

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/66, 31/41		A1	(11) International Publication Number: WO 98/13046 (43) International Publication Date: 2 April 1998 (02.04.98)															
(21) International Application Number: PCT/US97/14344 (22) International Filing Date: 15 August 1997 (15.08.97) (30) Priority Data: <table border="0"><tr><td>08/718,703</td><td>27 September 1996 (27.09.96)</td><td>US</td></tr><tr><td>08/842,360</td><td>24 April 1997 (24.04.97)</td><td>US</td></tr><tr><td>08/858,985</td><td>27 May 1997 (27.05.97)</td><td>US</td></tr><tr><td>08/863,624</td><td>27 May 1997 (27.05.97)</td><td>US</td></tr><tr><td>08/884,479</td><td>27 June 1997 (27.06.97)</td><td>US</td></tr></table> (71) Applicant: GUILFORD PHARMACEUTICALS INC. [US/US]; 611 Tributary Street, Baltimore, MD 21224 (US). (72) Inventors: SLUSHER, Barbara, S.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). JACKSON, Paul, F.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). TAYS, Kevin, L.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). MACLIN, Keith, M.; Guilford Pharmaceuticals Inc., 6611 Tributary Street, Baltimore, MD 21224 (US). (74) Agent: NATH, Gary, M.; Nath & Associates, 1835 K Street, N.W. #750, Washington, DC 20006-1203 (US).		08/718,703	27 September 1996 (27.09.96)	US	08/842,360	24 April 1997 (24.04.97)	US	08/858,985	27 May 1997 (27.05.97)	US	08/863,624	27 May 1997 (27.05.97)	US	08/884,479	27 June 1997 (27.06.97)	US	(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
08/718,703	27 September 1996 (27.09.96)	US																
08/842,360	24 April 1997 (24.04.97)	US																
08/858,985	27 May 1997 (27.05.97)	US																
08/863,624	27 May 1997 (27.05.97)	US																
08/884,479	27 June 1997 (27.06.97)	US																
(54) Title: NAALADASE COMPOSITIONS AND METHODS FOR TREATING GLUTAMATE ABNORMALITY AND EFFECTING NEURONAL ACTIVITY IN ANIMALS (57) Abstract <p>The present invention relates to a method of treating a glutamate abnormality and a method of effecting a neuronal activity in an animal using a NAALADase inhibitor, and a pharmaceutical composition comprising an effective amount of a NAALADase inhibitor for treating a glutamate abnormality and effecting a neuronal activity in an animal.</p>																		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauretania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NI	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

NAALADASE COMPOSITIONS AND METHODS FOR TREATING GLUTAMATE ABNORMALITY AND EFFECTING NEURONAL ACTIVITY IN ANIMALS

5 This application is a continuation-in-part of U.S. Patent Application No. 08/718,703, filed September 27, 1996, entitled "TREATMENT OF GLOBAL AND FOCAL ISCHEMIA USING NAALADASE INHIBITORS"; U.S. Patent Application No. 08/842,360 filed April 24, 1997, entitled "PHOSPHONIC
10 ACID DERIVATIVES"; U.S. Patent Application filed May 27, 1997, Attorney Docket No. 23029-X (serial number not yet assigned), entitled "NAALADASE INHIBITORS"; and U.S. Patent Application filed May 27, 1997, Attorney Docket No. 23029-X2 (serial number not yet assigned), entitled
15 "NAALADASE INHIBITORS", the entire contents of which applications are herein incorporated by reference.

BACKGROUND OF THE INVENTION

20 1. Field of the Invention

 The present invention relates to a method of treating a glutamate abnormality and a method of effecting a neuronal activity in an animal using a NAALADase inhibitor, and a pharmaceutical composition
25 comprising an effective amount of a NAALADase inhibitor for treating a glutamate abnormality and effecting a neuronal activity in an animal.

2. Description of the Prior Art

Glutamate Abnormalities

Glutamate has been implicated in various neurological diseases and conditions, including epilepsy, stroke, Alzheimer's disease, Parkinson's Disease, Amyotrophic Lateral Sclerosis (ALS), Huntington's Disease, schizophrenia, chronic pain, ischemia and neuronal loss following hypoxia, hypoglycemia, ischemia, trauma and nervous insult. Neurons release glutamate in great quantities when they are deprived of oxygen, as may occur during an ischemic brain insult such as a stroke or a heart attack. This excess release of glutamate in turn causes over-stimulation (excitotoxicity) of NMDA, AMPA, Kainate and MGR receptors. When glutamate binds to these receptors, ion channels in the cell membranes of the neurons open, permitting flows of ions across the cell membranes, e.g., Ca^{2+} and Na^+ into the cells and K^+ out of the cells. These flows of ions, especially the influx of Ca^{2+} , cause over-stimulation of the neurons. The over-stimulated neurons secrete more glutamate, creating a domino-effect which ultimately results in cell death via the production of proteases, lipases and free radicals.

Attempts to prevent excitotoxicity by blocking NMDA, AMPA, Kainate and MGR receptors have proven difficult because each receptor has multiple sites to which glutamate may bind. Many of the compositions that are

effective in blocking the receptors are also toxic to animals. As such, there is currently no known effective treatment for glutamate abnormalities.

NAALADase Inhibitors

5 NAAG and NAALADase have been implicated in several human and animal pathological conditions. For example, it has been demonstrated that intra-hippocampal injections of NAAG elicit prolonged seizure activity. More recently, it was reported that rats genetically
10 prone to epileptic seizures have a persistent increase in their basal level of NAALADase activity. These observations support the hypothesis that increased availability of synaptic glutamate elevates seizure susceptibility, and suggest that NAALADase inhibitors may
15 provide anti-epileptic activity.

 NAAG and NAALADase have also been implicated in the pathogenesis of ALS and in the pathologically similar animal disease called Hereditary Canine Spinal Muscular Atrophy (HCSMA). It has been shown that concentrations
20 of NAAG and its metabolites -- NAA, glutamate and aspartate -- are elevated two- to three-fold in the cerebrospinal fluid of ALS patients and HCSMA dogs. Additionally, NAALADase activity is significantly increased (two- to three-fold) in post-mortem spinal cord
25 tissue from ALS patients and HCSMA dogs. As such, NAALADase inhibitors may be clinically useful in curbing

the progression of ALS if increased metabolism of NAAG is responsible for the alterations of CSF levels of these acidic amino acids and peptides.

Abnormalities in NAAG levels and NAALADase activity
5 have also been documented in post-mortem schizophrenic brain, specifically in the prefrontal and limbic brain regions.

The findings described above suggest that NAALADase inhibitors could be useful in treating glutamate
10 abnormalities. In fact, the results of studies conducted by the inventors confirm that NAALADase inhibitors are effective in treating glutamate abnormalities, particularly stroke, Parkinson's Disease, Amyotrophic Lateral Sclerosis (ALS) and spinal cord injury.

15 While a few NAALADase inhibitors have been identified, they have only been used in non-clinical research. Examples of such inhibitors include general metallopeptidase inhibitors such as α -phenanthroline, metal chelators such as EGTA and EDTA, and peptide
20 analogs such as quisqualic acid and β -NAAG. Accordingly, a need exists for new NAALADase inhibitors, as well as pharmaceutical compositions and methods using such new and known NAALADase inhibitors to treat glutamate abnormalities.

25

SUMMARY OF THE INVENTION

The present invention relates to a pharmaceutical composition comprising:

- (i) an effective amount of a NAALADase inhibitor
5 for treating a glutamate abnormality or effecting a neuronal activity in an animal; and
- (ii) a pharmaceutically acceptable carrier.

The present invention further relates to a method of
10 treating a glutamate abnormality in an animal, comprising administering an effective amount of a NAALADase inhibitor to said animal.

The present invention also relates to a method of effecting a neuronal activity in an animal, comprising
15 administering an effective amount of a NAALADase inhibitor to said animal.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a bar graph plotting in vitro toxicity of
20 ischemic insult (potassium cyanide and 2-deoxyglucose) against various doses of 2-(phosphonomethyl)pentanedioic acid with which cortical cell cultures were treated.

FIG. 2 is a bar graph plotting in vitro toxicity against various doses of NAAG to which cortical cell
25 cultures were exposed.

FIG. 3 is a bar graph plotting in vitro toxicity following treatment with 2-(phosphonomethyl)pentanedioic acid, against various doses of NAAG to which cortical cell cultures were exposed.

5 FIG. 4 is a bar graph plotting in vitro toxicity of ischemic insult against various times at which cortical cell cultures were treated with 2-(phosphonomethyl)-pentanedioic acid.

FIG. 5 is a bar graph plotting in vivo cortical
10 injury volume against various doses of 2-(phosphonomethyl)pentanedioic acid with which rats were treated after sustaining middle cerebral artery occlusion.

FIG. 6 is a bar graph plotting in vivo total brain
15 infarct volume of rats against various times at which the rats are treated with 2-(phosphonomethyl)pentanedioic acid after sustaining middle cerebral artery occlusion.

FIG. 7 is a bar graph plotting in vivo extracellular glutamate increases in the striatum of rats against
20 various times at which the rats are treated with a vehicle or 2-(phosphonomethyl)pentanedioic acid after sustaining middle cerebral artery occlusion.

FIG. 8 is a bar graph plotting in vivo extracellular glutamate increases in the parietal cortex of rats
25 against various times at which the rats are treated with a vehicle or 2-(phosphonomethyl)pentanedioic acid after

sustaining middle cerebral artery occlusion.

FIG. 9 is a bar graph plotting in vivo extracellular glutamate increases in the frontal cortex of rats against various times at which the rats are treated with a vehicle or 2-(phosphonomethyl)pentanedioic acid after sustaining middle cerebral artery occlusion.

FIG. 10(a) is a photomicrograph of mouse sciatic nerve treated with a vehicle following cryolesion.

FIG. 10(b) is a photomicrograph of mouse sciatic nerve treated with 2-(phosphonomethyl)pentanedioic acid following cryolesion.

FIG. 11 is a bar graph plotting percent striatal TH innervation density against the treatment of mice with vehicle alone, vehicle following MPTP, or 2-(phosphonomethyl)pentanedioic acid following MPTP.

FIG. 12 is a bar graph plotting the neurological function code against the treatment of rats with dynorphin A alone or 2-(phosphonomethyl)pentanedioic acid with dynorphin A.

FIG. 13 is a bar graph plotting the ChAT activity of rat spinal cord organotypic cultures against the treatment of the cultures with 2-(phosphonomethyl)pentanedioic acid alone, THA alone, or THA with 2-(phosphonomethyl)pentanedioic acid.

FIG. 14 is a bar graph plotting the ChAT activity of rat spinal cord organotypic cultures against various

doses of 2-(phosphonomethyl)pentanedioic acid with which the cultures were treated in the presence of THA.

DETAILED DESCRIPTION OF THE INVENTION

5

Definitions

"Compound 3" refers to 2-(phosphonomethyl)pentanedioic acid (PMPA).

"Glutamate abnormality" refers to any disease, disorder or condition in which glutamate is implicated, including pathological conditions involving elevated levels of glutamate. Examples of glutamate abnormalities include epilepsy, stroke, Alzheimer's disease, Parkinson's Disease, Amyotrophic Lateral Sclerosis (ALS), Huntington's Disease, schizophrenia, chronic pain, ischemia and neuronal insult.

"Glutamate modulator" refers to any composition of matter which alone or in combination with another agent affects the level of glutamate in an animal.

"Inhibition", in the context of enzymes, refers to reversible enzyme inhibition such as competitive, uncompetitive and non-competitive inhibition. Competitive, uncompetitive and non-competitive inhibition can be distinguished by the effects of an inhibitor on the reaction kinetics of an enzyme. Competitive inhibition occurs when the inhibitor combines reversibly with the enzyme in such a way that it competes with a

normal substrate for binding at the active site. The affinity between the inhibitor and the enzyme may be measured by the inhibitor constant, K_i , which is defined as:

5

$$K_i = \frac{[E][I]}{[EI]}$$

10 wherein [E] is the concentration of the enzyme, [I] is the concentration of the inhibitor, and [EI] is the concentration of the enzyme-inhibitor complex formed by the reaction of the enzyme with the inhibitor. Unless otherwise specified, K_i as used herein refers to the
15 affinity between the inventive compounds and NAALADase. "IC₅₀" is a related term used to define the concentration or amount of a compound which is required to cause a 50% inhibition of the target enzyme.

"Ischemia" refers to localized tissue anemia due to
20 obstruction of the inflow of arterial blood. Global ischemia occurs when blood flow to the entire brain ceases for a period of time, such as may result from cardiac arrest. Focal ischemia occurs when a portion of the brain is deprived of its normal blood supply, such as
25 may result from thromboembolytic occlusion of a cerebral vessel, traumatic head injury, edema or brain tumor. Even if transient, both global and focal ischemia can produce widespread neuronal damage. Although nerve

10


tissue damage occurs over hours or even days following the onset of ischemia, some permanent nerve tissue damage may develop in the initial minutes following cessation of blood flow to the brain. Much of this damage is attributed to glutamate toxicity and secondary consequences of reperfusion of the tissue, such as the release of vasoactive products by damaged endothelium, and the release of cytotoxic products, such as free radicals and leukotrienes, by the damaged tissue.

5 "NAAG" refers to N-acetyl-aspartyl-glutamate, an important peptide component of the brain, with levels comparable to the major inhibitor neurotransmitter gamma-aminobutyric acid (GABA). NAAG is neuron-specific, present in synaptic vesicles and released upon neuronal stimulation in several systems presumed to be glutamatergic. Studies suggest that NAAG may function as a neurotransmitter and/or neuromodulator in the central nervous system, or as a precursor of the neurotransmitter glutamate.

15 "NAALADase" refers to N-acetylated α -linked acidic dipeptidase, a membrane-bound metallopeptidase which catabolizes NAAG to N-acetylaspartate (NAA) and glutamate:

25

Catabolism of NAAG by NAALADase



15 NAALADase shows a high affinity for NAAG with a K_m of 540 nM. If NAAG is a bioactive peptide, then NAALADase may serve to inactivate NAAG'S synaptic action. Alternatively, if NAAG functions as a precursor for glutamate, the primary function of NAALADase may be to

20 regulate synaptic glutamate availability.

"Nervous function" refers to the various functions of the nervous system, which among other things provide an awareness of the internal and external environments of the body, make possible voluntary and reflex activities

25 between the various structural elements of the organism, and balance the organism's response to environmental changes.

"Nervous insult" refers to any damage to nervous tissue and any disability or death resulting therefrom.

30 The cause of nervous insult may be metabolic, toxic, neurotoxic, iatrogenic, thermal or chemical, and includes

without limitation ischemia, hypoxia, cerebrovascular accident, trauma, surgery, pressure, mass effect, hemorrhage, radiation, vasospasm, neurodegenerative disease, neurodegenerative process, infection,
5 Parkinson's disease, ALS, myelination/demyelination process, epilepsy, cognitive disorder, glutamate abnormality and secondary effects thereof. Currently, there is no known effective treatment for nervous tissue damage.

10 "Nervous tissue" refers to the various components that make up the nervous system, including without limitation neurons, neural support cells, glia, Schwann cells, vasculature contained within and supplying these structures, the central nervous system, the brain, the
15 brain stem, the spinal cord, the junction of the central nervous system with the peripheral nervous system, the peripheral nervous system and allied structures.

"Neuroprotective" refers to the effect of reducing, arresting or ameliorating nervous insult, and protecting,
20 resuscitating or reviving nervous tissue which has suffered nervous insult.

"Pharmaceutically acceptable salt" refers to a salt of the inventive compounds which possesses the desired pharmacological activity and which is neither
25 biologically nor otherwise undesirable. The salt can be formed with inorganic acids such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate butyrate, citrate, camphorate,

camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, thiocyanate, tosylate and undecanoate. Examples of a base salt include ammonium salts, alkali metal salts such as sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine and lysine. The basic nitrogen-containing groups can be quarternized with agents including lower alkyl halides such as methyl, ethyl, propyl and butyl chlorides, bromides and iodides; dialkyl sulfates such as dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; and aralkyl halides such as benzyl and phenethyl bromides.

"Treatment" refers to:

- (i) preventing a disease, disorder or condition from occurring in an animal which may be predisposed to the disease, disorder and/or condition but has not yet been diagnosed as having it;

14

(ii) inhibiting the disease, disorder or condition, i.e., arresting its development; and

(iii) relieving the disease, disorder or condition, i.e., causing regression of the disease, disorder and/or
5 condition.

PHARMACEUTICAL COMPOSITIONS OF THE PRESENT INVENTION

The present invention relates to a pharmaceutical composition comprising:

- 10 (i) an effective amount of a NAALADase inhibitor for treating a glutamate abnormality or effecting a neuronal activity in an animal; and
- (ii) a pharmaceutically acceptable carrier.

15 The pharmaceutical composition may further comprise at least one additional therapeutic agent.

Since NAALADase is a metallopeptidase, useful NAALADase inhibitors for the pharmaceutical composition of the present invention include small molecule compounds
20 with functional groups known to inhibit metallo-peptidases, such as hydroxyphosphinyl derivatives.


According to scientific literature, the glutamate moiety plays a more critical role than the aspartate moiety in the recognition of NAAG by NAALADase. As such,
25 a preferred NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative, an acidic peptide analog or

15

a mixture thereof.

A preferred acidic peptide analog is selected from the group consisting of Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-Glu and Glu-Glu-Glu.

5 A preferred NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative of formula I:



10

or a pharmaceutically acceptable salt or hydrate thereof, wherein:

R_1 is selected from the group consisting of
 15 hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl and Ar, wherein said R₁ is unsubstituted or substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 -
 20 C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_9 alkoxy, C_2 - C_9 alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

X is CR_3R_4 , O or NR_1 ;

R_3 and R_4 are independently selected from the group
 25 consisting of hydrogen, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_3 - C_8

cycloalkyl, C₅-C₇ cycloalkenyl, Ar, halo and mixtures thereof;

R₂ is selected from the group consisting of hydrogen, C₁-C₆ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl and Ar, wherein said R₂ is unsubstituted or substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

Ar is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, benzyl and phenyl, wherein said Ar is unsubstituted or substituted with halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino or mixtures thereof.

Preferably, X is CH₂.

More preferably, R₂ is substituted with carboxy.

Even more preferably, R₁ is hydrogen, C₁-C₄ straight or branched chain alkyl, C₂-C₄ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl or phenyl, wherein said R₁ is unsubstituted or substituted

with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, 5 benzyloxy, amino, benzyl, phenyl or mixtures thereof; and R₂ is C₁-C₂ alkyl.

Most preferably, the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 10 2-(phosphonomethyl)pentanedioic acid;
- 2-(phosphonomethyl)succinic acid;
- 2-[[(2-carboxyethyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(cyclohexyl)methylhydroxyphosphinyl]methyl]pentane-
- 20 dioic acid;
- 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[benzylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(phenylmethyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 25 2-[[(phenylethyl)hydroxyphosphinyl]methyl]pentanedioic acid;

- 2-[[(phenylpropyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(phenylbutyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 5 2-[[(4-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(4-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(2-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic
- 10 acid;
- 2-[[(pentafluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(methoxybenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 15 2-[[(2,3,4-trimethoxyphenyl) hydroxyphosphinyl] methyl] - pentanedioic acid;
- 2-[[(phenylprop-2-enyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(2-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic
- 20 acid;
- 2-[[((hydroxy) phenylmethyl) hydroxyphosphinyl] methyl] - pentanedioic acid;
- 2-[[(3-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 25 2-[[(4-fluorophenyl) hydroxyphosphinyl] methyl] pentanedioic acid;

2-[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]-
pentanedioic acid; and

pharmaceutically acceptable salts and hydrates thereof.

In other embodiments, R_2 is C_3 - C_9 alkyl; R_1 is 2-
5 indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl,
tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-
thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl or C_1 - C_4
straight or branched chain alkyl substituted with 2-
indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl,
10 tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-
pyridyl or 4-pyridyl; or R_1 is 1-naphthyl, 2-naphthyl, or
 C_1 - C_4 straight or branched chain alkyl substituted with
1-naphthyl or 2-naphthyl.

Preferred compounds of these embodiments include:

- 15 2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
20 2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]decanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid;
25 2-[[(2-pyridyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;

- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 5 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]methyl]-
10 pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]methyl]-
pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]methyl]-
pentanedioic acid;
- 15 2-[[[(2-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
pentanedioic acid;
- 2-[[[(3-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
pentanedioic acid;
- 2-[[[(4-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
20 pentanedioic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 25 2-[[[(4-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;

- 2-[[[(3-indolyl)ethylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 5 2-[[[(2-thienyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl)methyl]pentane-
10 dioic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 15 2-[[[(2-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 2-[[[(3-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 2-[[[(4-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
20 acid;
- 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[[(2-indolyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 25 2-[[[(3-indolyl)hydroxyphosphinyl)methyl]pentanedioic
acid;

- 2-[[[(4-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 5 2-[[[(3-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(4-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(1-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2-[[[(2-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(1-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2-[[[(2-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 20 2-[[[(1-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 25 2-[[[(1-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic acid;

23.

2-[[(2-naphthyl)butylhydroxyphosphinyl]methyl]pentane-dioic acid; and

pharmaceutically acceptable salts and hydrates thereof.

In another preferred embodiment, X is CH₂ and R₂ is
5 selected from the group consisting of hydrogen, C₁-C₆,
straight or branched chain alkyl, C₂-C₆, straight or
branched chain alkenyl, C₃-C₈, cycloalkyl, C₅-C₇,
cycloalkenyl, benzyl and phenyl, wherein said R₂ is
unsubstituted or substituted with C₃-C₈, cycloalkyl, C₅-C₇,
10 cycloalkenyl, C₁-C₆, straight or branched chain alkyl, C₂-
C₆, straight or branched chain alkenyl, C₁-C₄, alkoxy,
phenyl or mixtures thereof.

More preferably, R₁ is hydrogen, C₁-C₆, straight or
branched chain alkyl, C₂-C₆, straight or branched chain
15 alkenyl, C₃-C₈, cycloalkyl, C₅-C₇, cycloalkenyl, benzyl or
phenyl, wherein said R₁ is unsubstituted or substituted
with carboxy, C₃-C₈, cycloalkyl, C₅-C₇, cycloalkenyl, halo,
hydroxy, nitro, trifluoromethyl, C₁-C₆, straight or
branched chain alkyl, C₂-C₆, straight or branched chain
20 alkenyl, C₁-C₄, alkoxy, C₂-C₄, alkenyloxy, phenoxy,
benzyloxy, amino, benzyl, phenyl or mixtures thereof.

Most preferably, the glutamate-derived
hydroxyphosphinyl derivative is selected from the group
consisting of:

- 25 3-(methylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(ethylhydroxyphosphinyl)-2-phenylpropanoic acid;

- 3- (propylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- (butylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- (cyclohexylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- ((cyclohexyl)methylhydroxyphosphinyl) -2-phenylpropanoic
5 acid;
3- (phenylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- (phenylethylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- (phenylpropylhydroxyphosphinyl) -2-phenylpropanoic acid;
10 3- (phenylbutylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- ((2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl) -2-phenylpropanoic acid;
3- (phenylprop-2-enylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2-ethylpropanoic acid;
15 3- (benzylhydroxyphosphinyl) -2-propylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2-butylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2-cyclohexylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2- (cyclohexyl)methylpropanoic
20 acid;
3- (benzylhydroxyphosphinyl) -2-phenylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2-benzylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2-phenylethylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2-phenylpropylpropanoic acid;
25 3- (benzylhydroxyphosphinyl) -2-phenylbutylpropanoic acid;
3- (benzylhydroxyphosphinyl) -2- (2,3,4-trimethoxyphenyl) -

propanoic acid;

3-(benzylhydroxyphosphinyl)-2-phenylprop-2-enylpropanoic acid; and

pharmaceutically acceptable salts and hydrates thereof.

5 In other embodiments, at least one of R₁ and R₂ is 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or C₁-C₄ straight or branched chain alkyl substituted with 2-
10 indolyl 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl or 4-pyridyl; or R₁ is 1-naphthyl, 2-naphthyl, or C₁-C₄ straight or branched chain alkyl substituted with 1-naphthyl or 2-naphthyl.

15 Preferred compounds of these embodiments include:
3-[(2-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
20 3-[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
3-[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic
25 acid;
3-[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenyl

- propanoic acid;
3-[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenyl
propanoic acid;
3-[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenyl
5 propanoic acid;
3-[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
10 3-[(4-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-indolyl)ethylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-indolyl)propylhydroxyphosphinyl]-2-phenylpropanoic
15 acid;
3-[(2-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
20 3-[(4-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-thienyl)ethylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-thienyl)propylhydroxyphosphinyl]-2-phenylpropanoic
25 acid;
3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)methylpropanoic

- acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)methylpropanoic acid;
- 5 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propylpropanoic acid;
- 10 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propylpropanoic acid;
- 15 3-(benzylhydroxyphosphinyl)-2-(2-indolyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)methylpropanoic acid;
- 20 3-(benzylhydroxyphosphinyl)-2-(4-indolyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)propylpropanoic acid;
- 25 3-(benzylhydroxyphosphinyl)-2-(2-thienyl)methylpropanoic

- acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-thienyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(4-thienyl)methylpropanoic acid;
- 5 acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-thienyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-thienyl)propylpropanoic acid;
- 10 3-((1-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((1-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 15 acid;
- 3-((1-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 20 3-((1-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((1-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 25 acid;
- 3-((2-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic

acid; and

pharmaceutically acceptable salts and hydrates thereof.

When X is O, R₂ is preferably substituted with carboxy.

- 5 Exemplary compounds of this embodiment include:
- 2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;
10 2-[[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;
15 2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
20 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy] pentane-
25 dioic acid;
2-[[[(methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic

- acid;
- 2- [[(2,3,4-trimethoxyphenyl)hydroxyphosphinyl]oxy] -
pentanedioic acid;
- 2- [[(1-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 5 2- [[(2-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2- [[(1-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 2- [[(2-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 10 2- [[(1-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 2- [[(2-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 2- [[(1-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic
15 acid;
- 2- [[(2-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 2- [[(1-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 20 2- [[(2-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic
acid;
- 2- [[(phenylprop-2-enyl)hydroxyphosphinyl]oxy]pentanedioic
acid;
- 2- [[(benzylhydroxyphosphinyl]oxy]pentanedioic acid;
- 25 2- [[((hydroxy)phenylmethyl)hydroxyphosphinyl]oxy]pentane-
dioic acid;

- 2-[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 5 2-[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-(phosphono)oxy]pentanedioic acid;
- 2-[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]-pentanedioic acid;
- 10 2-[[methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[cyclohexylhydroxyphosphinyl]oxy]-2-phenylethanoic
- 15 acid;
- 2-[[(cyclohexyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 20 2-[[phenylethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic
- 25 acid;
- 2-[[(2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl]oxy]-2-

- phenylethanoic acid;
- 2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 5 2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 10 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 15 2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 20 2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;
- 25 2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;

- 2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;
5 2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]decanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-methylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-ethylethanoic acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-propylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-butylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-cyclohexylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(cyclohexyl)methyl-
15 ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-benzylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethylethanoic
acid;
20 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylpropylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylbutylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2,3,4-trimethoxy-
25 phenyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethanoic

- acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)methyl-
5 ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethyl-
ethanoic acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propyl-
15 ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butyl-
ethanoic acid;
20 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enyl-
ethanoic acid;
2-[[2-(pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[3-(pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic
25 acid;
2-[[4-(pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic

- acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 5 acid;
- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 10 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 15 acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 20 2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 25 acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic

acid;

2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;

2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic
5 acid; and

pharmaceutically acceptable salts and hydrates thereof.

In another preferred embodiment, R₂ is selected from
the group consisting of hydrogen, C₁-C₉ straight or
branched chain alkyl, C₂-C₉ straight or branched chain
10 alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl and
phenyl, wherein said R₂ is unsubstituted or substituted
with C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, C₁-C₆ straight
or branched chain alkyl, C₂-C₆ straight or branched chain
alkenyl, C₁-C₄ alkoxy, phenyl or mixtures thereof.

15 Exemplary compounds of this embodiment include:

2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-
ethanoic acid;

2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-
ethanoic acid;

20 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-
ethanoic acid;

2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-
ethanoic acid;

2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenyl-
25 ethanoic acid;

2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-

- phenylethanoic acid;
2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
5 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
10 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
15 2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
20 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
25 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)methyl-

- ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)methyl-ethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)propyl-ethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-propylethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)methyl-ethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)propyl-ethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)methyl-

- ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)methyl-
5 ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)propyl-
ethanoic acid; and
10 pharmaceutically acceptable salts and hydrates thereof.

When X is NR₁, R₂ is preferably substituted with carboxy.

Exemplary compounds of this embodiment include:

- 2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;
15 2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[cyclohexyl)methylhydroxyphosphinyl]amino]pentane-
20 dioic acid;
2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic
25 acid;
2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;

- 2-[[(4-methylbenzyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2-[[(4-fluorobenzyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 5 2-[[(2-fluorobenzyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2-[[(pentafluorobenzyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2-[[(methoxybenzyl) hydroxyphosphinyl] amino] pentanedioic
- 10 acid;
- 2-[[(2,3,4-trimethoxyphenyl) hydroxyphosphinyl] amino] - pentanedioic acid;
- 2-[[(1-naphthyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 15 2-[[(2-naphthyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2-[[(1-naphthyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 2-[[(2-naphthyl) methylhydroxyphosphinyl] amino] pentane-
- 20 dioic acid;
- 2-[[(1-naphthyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 2-[[(2-naphthyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 25 2-[[(1-naphthyl) propylhydroxyphosphinyl] amino] pentanedioic acid;

- 2-[[(2-naphthyl)propylhydroxyphosphinyl]amino]pentane-
dioic acid;
- 2-[[(1-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic
acid;
- 5 2-[[(2-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic
acid;
- 2-[[(phenylprop-2-enyl)hydroxyphosphinyl]amino]pentane-
dioic acid;
- 2-[[(benzylhydroxyphosphinyl)amino]pentanedioic acid;
- 10 2-[[(2-fluorobenzyl)hydroxyphosphinyl]amino]-2-pentane-
dioic acid;
- 2-[[((hydroxy)phenylmethyl)hydroxyphosphinyl]amino]-
pentanedioic acid;
- 2-[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic
15 acid;
- 2-[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic
acid;
- 2-[(phosphono)amino]pentanedioic acid;
- 2-[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]-
20 pentanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]heptanedioic acid;
- 25 2-[(methylhydroxyphosphinyl)amino]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]octanedioic acid;

- 2 - [(methylhydroxyphosphinyl) amino] nonanedioic acid;
2 - [(benzylhydroxyphosphinyl) amino] nonanedioic acid;
2 - [(methylhydroxyphosphinyl) amino] decanedioic acid;
2 - [(benzylhydroxyphosphinyl) amino] decanedioic acid;
5 3 - [[(2-pyridyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
3 - [[(3-pyridyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
3 - [[(4-pyridyl) methylhydroxyphosphinyl] amino] pentanedioic
10 acid;
3 - [[(3-pyridyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
3 - [[(3-pyridyl) propylhydroxyphosphinyl] amino] pentanedioic acid;
15 3 - [[(tetrahydrofuranyl) methylhydroxyphosphinyl] amino] - pentanedioic acid;
3 - [[(tetrahydrofuranyl) ethylhydroxyphosphinyl] amino] - pentanedioic acid;
3 - [[(tetrahydrofuranyl) propylhydroxyphosphinyl] amino] -
20 pentanedioic acid;
3 - [[(2-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
3 - [[(3-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
25 3 - [[(4-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;

- 3- [[[3-indolyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 3- [[[3-indolyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 3- [[[2-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3- [[[3-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3- [[[4-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
- 10 acid;
- 3- [[[3-thienyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 3- [[[3-thienyl)propylhydroxyphosphinyl]amino]pentanedioic acid; and
- 15 pharmaceutically acceptable salts and hydrates thereof.

In another preferred embodiment, R_2 is selected from the group consisting of hydrogen, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_3 - C_6 cycloalkyl, C_5 - C_6 cycloalkenyl, benzyl and

20 phenyl, wherein said R_2 is unsubstituted or substituted with C_3 - C_6 cycloalkyl, C_5 - C_6 cycloalkenyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_6 alkoxy, phenyl or mixtures thereof.

Exemplary compounds of this embodiment include:

- 25 2- [[methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2- [[ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

- 2-[[propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[cyclohexylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
5 2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
2-[[phenylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[phenylethylhydroxyphosphinyl]amino]-2-phenylethanoic
10 acid;
2-[[phenylpropylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[phenylbutylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
15 2-[[[(2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic
20 acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
25 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;

- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 5 2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 10 2-[[[phenylprop-2-enylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-ethylethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-butylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-cyclohexylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(cyclohexyl)methyl-ethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-benzylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]amino]-2-phenylpropylethanoic acid;

- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylbutylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2,3,4-trimethoxyphenyl)ethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)methyl-
- 10 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethyl-ethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)propyl-
- 20 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)butyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)butyl-ethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]amino]-2-phenolprop-2-enyl-ethanoic acid;

- 2-[[(2-pyridyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[(3-pyridyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 5 2-[[(4-pyridyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[(3-pyridyl)ethylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[(3-pyridyl)propylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 10 2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 15 2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
- 2-[[(2-indolyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[(3-indolyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 20 2-[[(4-indolyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[(3-indolyl)ethylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 25 2-[[(3-indolyl)propylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;


- 2-[[[(2-thienyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 5 2-[[[(4-thienyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)methyl-ethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)propyl-ethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)-methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)-ethylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)-propylethanoic acid;

- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)methyl-ethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)propyl-10 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)methyl-ethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)propyl-20 ethanoic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

Synthesis of NAALADase Inhibitors

The NAALADase inhibitors of formula I can be readily
25 prepared by standard techniques of organic chemistry,
utilizing the general synthetic pathways depicted below

in Schemes I-IX. Precursor compounds can be prepared by methods known in the art, such as those described by Jackson et al., J. Med. Chem., Vol. 39, No. 2, pp. 619-622 (1996) and Froestl et al., J. Med. Chem., Vol. 38, pp. 3313-3331 (1995).


Scheme I

Methods of substituting the R group are known in the art. Additional methods of synthesizing phosphinic acid esters are described in J. Med. Chem., Vol. 31, pp. 204-212 (1988), and set forth below in Scheme II.

51


Scheme II

Method A




|||

10



- | | | | |
|-------|--|----|---|
| A. | $\text{R}' = (\text{CH}_2)_3\text{Ph}$ | H. | $\text{R}' = n\text{-C}_7\text{H}_{15}$ |
| 15 B. | $(\text{CH}_2)_4\text{Ph}$ | I. | $n\text{-C}_8\text{H}_{17}$ |
| C. | $(\text{CH}_2)_5\text{Ph}$ | J. | $n\text{-C}_9\text{H}_{19}$ |
| D. | $(\text{CH}_2)_4(\text{P-F-Ph})$ | K. | $\text{CH}_2\text{CHCH}_3\text{C}_4\text{H}_9$ |
| E. | $(\text{CH}_2)_4-(3\text{-pyridyl})$ | L. | $\text{CH}_2(\text{CH}_3)\text{C}(\text{CH}_3)_2$ |
| F. | $n\text{-C}_5\text{H}_{11}$ | | |
| 20 G. | $n\text{-C}_6\text{H}_{13}$ | | |

Method B




52

- N. $R' = n-C_4H_9$
 O. $CHCH_3C_5H_{11}$

Starting with the aforementioned phosphinic acid
 5 esters, there are a variety of routes for preparing the
 compounds of formula I. For example, a general route has
 been described in J. Med. Chem., Vol. 39, pp. 619-622
 (1996), and is set forth below in Scheme III.

10


Scheme III

Other routes for preparing the compounds of formula
 I are set forth below in Scheme IV and Scheme V. Scheme
 25 IV and Scheme V show the starting material as a
 phosphinic acid derivative and the R group as any


53

reasonable chemical substituent including without limitation the substituents listed in Scheme II and throughout the specification.

5


Scheme IV

15

Scheme V


54

Another route for preparing the compounds of formula I allows for aromatic substitution at R_1 , and is set forth below in Scheme VI.




Another route for preparing the compounds of formula I allows for aromatic substitution at the R_2 position, and is set forth below in Scheme VII.

55

Scheme VII


Another route for preparing the compounds of formula I wherein X is NR₁ is set forth below in Scheme VIII.

Scheme VIII



Another route for preparing the compounds of formula I wherein X is oxygen is set forth below in Scheme IX.

57

Scheme IXMETHODS OF THE PRESENT INVENTIONMETHODS OF TREATING A GLUTAMATE ABNORMALITY

15 Although not limited to any one particular theory, it is believed that the NAALADase inhibitors used in the methods of the present invention modulate levels of glutamate by acting on a storage form of glutamate which is hypothesized to be upstream from the effects mediated

20 by the NMDA receptor.

Accordingly, the present invention further relates to a method of treating a glutamate abnormality in an animal, comprising administering an effective amount of a NAALADase inhibitor to said animal.

25 The glutamate abnormality may be any disease, disorder or condition in which glutamate is implicated,


including pathological conditions involving elevated levels of glutamate. Examples of glutamate abnormalities include without limitation epilepsy, stroke, Alzheimer's disease, Parkinson's Disease, Amyotrophic Lateral Sclerosis (ALS), Huntington's Disease, schizophrenia, chronic pain, ischemia, peripheral neuropathy, traumatic brain injury and physical damage to the spinal cord. In a preferred embodiment, the glutamate abnormality is selected from the group consisting of stroke, Parkinson's Disease, Amyotrophic Lateral Sclerosis (ALS) and spinal cord injury.

The NAALADase inhibitor may be administered alone or in combination with at least one additional therapeutic agent.

A preferred NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative, an acidic peptide analog or a mixture thereof.

A preferred acidic peptide analog is an amino acid sequence selected from the group consisting of Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-Glu and Glu-Glu-Glu.

A preferred NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative of formula I:



I

or a pharmaceutically acceptable salt or hydrate thereof,
wherein:

R_1 is selected from the group consisting of hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl and Ar, wherein said R_1 is unsubstituted or substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

X is CR_3R_4 , O or NR_1 ;

R_3 and R_4 are independently selected from the group consisting of hydrogen, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, Ar, halo and mixtures thereof;

R_2 is selected from the group consisting of hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl and Ar, wherein said R_2 is unsubstituted or substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_6 alkoxy, C_2 - C_6 alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

Ar is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, benzyl and phenyl, wherein said Ar is unsubstituted or substituted with halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino or mixtures thereof.

10 Preferably, X is CH₂.

More preferably, R₂ is substituted with carboxy.

Even more preferably, R₁ is hydrogen, C₁-C₄ straight or branched chain alkyl, C₂-C₄ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl or phenyl, wherein said R₁ is unsubstituted or substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof; and R₂ is C₁-C₂ alkyl.

Most preferably, the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 25 2-(phosphonomethyl)pentanedioic acid;
2-(phosphonomethyl)succinic acid;

- 2-[[(2-carboxyethyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[methylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[ethylhydroxyphosphinyl)methyl]pentanedioic acid;
- 5 2-[[propylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[butylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[(cyclohexyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
- 10 2-[[phenylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[(phenylmethyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[(phenylethyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 15 2-[[(phenylpropyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[(phenylbutyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 20 2-[[(4-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[(4-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 25 2-[[(pentafluorobenzyl)hydroxyphosphinyl)methyl]pentane-

- dioic acid;
- 2-[[[(methoxybenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2,3,4-trimethoxyphenyl)hydroxyphosphinyl)methyl]-
- 5 pentanedioic acid;
- 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 10 2-[[[(hydroxy)phenylmethyl)hydroxyphosphinyl)methyl]-pentanedioic acid;
- 2-[[[(3-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic
- 15 acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]-pentanedioic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

Examples of other glutamate-derived

20 hydroxyphosphinyl derivatives useful for the present method are identified above with respect to pharmaceuticals compositions.

METHODS OF EFFECTING A NEURONAL ACTIVITY

- 25 The inventors have discovered that inhibition of NAALADase promotes nerve regeneration and myelin

formation.

Accordingly, the present invention also relates to a method of effecting a neuronal activity in an animal, comprising administering an effective amount of a
5 NAALADase inhibitor to said animal.

The neuronal activity that is effected by the method of the present invention may be selected from the group consisting of: stimulation of damaged neurons, promotion of neuronal regeneration, prevention of neurodegeneration
10 and treatment of a neurological disorder.

Examples of a neurological disorder that is treatable by the method of the present invention include without limitation: trigeminal neuralgia; glossopharyngeal neuralgia; Bell's Palsy; myasthenia
15 gravis; muscular dystrophy; amyotrophic lateral sclerosis; progressive muscular atrophy; progressive bulbar inherited muscular atrophy; herniated, ruptured or prolapsed invertebrate disk syndromes; cervical spondylosis; plexus disorders; thoracic outlet
20 destruction syndromes; peripheral neuropathies such as those caused by lead, dapsone, ticks, porphyria, or Guillain-Barré syndrome; Alzheimer's disease; and Parkinson's disease.

The method of the present invention is particularly
25 useful for treating a neurological disorder selected from the group consisting of: peripheral neuropathy caused by

physical injury or disease state, traumatic brain injury, physical damage to the spinal cord, stroke associated with brain damage, demyelinating diseases and neurological disorders relating to neurodegeneration.

5 Examples of demyelinating diseases include multiple sclerosis. Examples of neurological disorders relating to neurodegeneration include Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis (ALS).


10 The NAALADase inhibitor may be administered alone or in combination with at least one additional therapeutic agent.

A preferred NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative, an acidic peptide
15 analog or a mixture thereof.

A preferred acidic peptide analog is an amino acid sequence selected from the group consisting of Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-Glu and Glu-Glu-Glu.

A preferred NAALADase inhibitor is a glutamate-
20 derived hydroxyphosphinyl derivative of formula I:

65



5

or a pharmaceutically acceptable salt or hydrate thereof,
wherein:

R₁ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉
10 straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl and Ar, wherein said R₁ is unsubstituted or substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or
15 branched chain alkenyl, C₁-C₉ alkoxy, C₂-C₉ alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

X is CR₃R₄, O or NR₁;

R₃ and R₄ are independently selected from the group consisting of hydrogen, C₁-C₆ straight or branched chain
20 alkyl, C₂-C₆ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, Ar, halo and mixtures thereof;

R₂ is selected from the group consisting of hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉
25 straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl and Ar, wherein said R₂ is unsubstituted

or substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

Ar is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuran-2-yl, tetrahydropyran-2-yl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, benzyl and phenyl, wherein said Ar is unsubstituted or substituted with halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino or mixtures thereof.

Preferably, X is CH₂.

More preferably, R₂ is substituted with carboxy.

Even more preferably, R₁ is hydrogen, C₁-C₄ straight or branched chain alkyl, C₂-C₄ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl or phenyl, wherein said R₁ is unsubstituted or substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof; and R₂ is C₁-C₂ alkyl.

Most preferably, the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 2-(phosphonomethyl)pentanedioic acid;
- 5 2-(phosphonomethyl)succinic acid;
- 2-[[(2-carboxyethyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(cyclohexyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[(phenylmethyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(phenylethyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 20 2-[[(phenylpropyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[(phenylbutyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 25 2-[[(4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic acid;

- 2-[[[(4-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 5 2-[[[(pentafluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(methoxybenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2,3,4-trimethoxyphenyl)hydroxyphosphinyl)methyl]-
- 10 pentanedioic acid;
- 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 15 2-[[[(hydroxy)phenylmethyl)hydroxyphosphinyl)methyl]-pentanedioic acid;
- 2-[[[(3-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic
- 20 acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]-pentanedioic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

Examples of other glutamate-derived

25 hydroxyphosphinyl derivatives useful for the present method are identified above with respect to

pharmaceuticals compositions.

ROUTE OF ADMINISTRATION

In the methods of the present invention, the
5 compounds may be administered orally, parenterally, by
inhalation spray, topically, rectally, nasally, buccally,
vaginally or via an implanted reservoir in dosage
formulations containing conventional non-toxic
pharmaceutically-acceptable carriers, adjuvants and
10 vehicles. The term parenteral as used herein includes
subcutaneous, intravenous, intramuscular,
intraperitoneal, intrathecal, intraventricular,
intrasternal or intracranial injection and infusion
techniques. Invasive techniques are preferred,
15 particularly direct administration to damaged neuronal
tissue.

To be effective therapeutically as central nervous
system targets, the NAALADase inhibitors used in the
methods of the present invention should readily penetrate
20 the blood-brain barrier when peripherally administered.
Compounds which cannot penetrate the blood-brain barrier
can be effectively administered by an intraventricular
route.

The compounds may also be administered in the form
25 of sterile injectable preparations, for example, as
sterile injectable aqueous or oleaginous suspensions.

These suspensions can be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparations may also be sterile injectable solutions or suspensions in non-toxic parenterally-acceptable diluents or solvents, for example, as solutions in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

10 In addition, sterile fixed oils are conventionally employed as solvents or suspending mediums. For this purpose, any bland fixed oil such as a synthetic mono- or di-glyceride may be employed. Fatty acids such as oleic acid and its glyceride derivatives, including olive oil

15 and castor oil, especially in their polyoxyethylated forms, are useful in the preparation of injectables. These oil solutions or suspensions may also contain long-chain alcohol diluents or dispersants.

Additionally, the compounds may be administered orally in the form of capsules, tablets, aqueous suspensions or solutions. Tablets may contain carriers such as lactose and corn starch, and/or lubricating agents such as magnesium stearate. Capsules may contain diluents including lactose and dried corn starch.

25 Aqueous suspensions may contain emulsifying and suspending agents combined with the active ingredient.

The oral dosage forms may further contain sweetening and/or flavoring and/or coloring agents.

The compounds may further be administered rectally in the form of suppositories. These compositions can be prepared by mixing the drug with suitable non-irritating excipients which are solid at room temperature, but liquid at rectal temperature such that they will melt in the rectum to release the drug. Such excipients include cocoa butter, beeswax and polyethylene glycols.

Moreover, the compounds may be administered topically, especially when the conditions addressed for treatment involve areas or organs readily accessible by topical application, including neurological disorders of the eye, the skin or the lower intestinal tract.

For topical application to the eye, or ophthalmic use, the compounds can be formulated as micronized suspensions in isotonic, pH adjusted sterile saline or, preferably, as a solution in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, the compounds may be formulated into ointments, such as petrolatum.

For topical application to the skin, the compounds can be formulated into suitable ointments containing the compounds suspended or dissolved in, for example, mixtures with one or more of the following: mineral oil,

liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated into suitable lotions or creams containing the
5 active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl ester wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

10 Topical application to the lower intestinal tract can be effected in rectal suppository formulations (see above) or in suitable enema formulations.

The NAALADase inhibitors used in the methods of the present invention may be administered by a single dose,
15 multiple discrete doses or continuous infusion. Since the compounds are small, easily diffusible and relatively stable, they are well suited to continuous infusion. Pump means, particularly subcutaneous pump means, are preferred for continuous infusion.

20

DOSAGE

Dose levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred
25 levels being about 0.1 mg to about 1,000 mg. The specific dose level for any particular patient will vary

depending upon a variety of factors, including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the patient; the time of administration; the rate of excretion; drug
5 combination; the severity of the particular disease being treated; and the form of administration. Typically, in vitro dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models are also helpful. The considerations for
10 determining the proper dose levels are well known in the art.

In a preferred embodiment, the NAALADase inhibitors are administered in lyophilized form. In this case, 1 to 100 mg of a NAALADase inhibitor may be lyophilized in
15 individual vials, together with a carrier and a buffer, such as mannitol and sodium phosphate. The compound may be reconstituted in the vials with bacteriostatic water before administration.

In treating global ischemia, the NAALADase
20 inhibitors are preferably administered orally, rectally, parenterally or topically at least 1 to 6 times daily, and may follow an initial bolus dose of higher concentration.

As previously mentioned, the NAALADase inhibitors
25 used in the methods of the present invention may be administered in combination with one or more therapeutic

agents. Specific dose levels for these agents will depend upon considerations such as those identified above for the NAALADase inhibitors.

5

ADMINISTRATION REGIMEN

For the methods of the present invention, any administration regimen regulating the timing and sequence of drug delivery can be used and repeated as necessary to effect treatment. Such regimen may include pretreatment
10 and/or co-administration with additional therapeutic agents.

To maximize protection of nervous tissue from nervous insult, the NAALADase inhibitors should be administered to the affected cells as soon as possible.
15 In situations where nervous insult is anticipated, the compounds should be administered before the expected nervous insult. Such situations of increased likelihood of nervous insult include surgery (cartoid endarterectomy, cardiac, vascular, aortic, orthopedic);
20 endovascular procedures such as arterial catheterization (cartoid, vertebral, aortic, cardia, renal, spinal, Adamkiewicz); injections of embolic agents; coils or balloons for hemostasis; interruptions of vascularity for treatment of brain lesions; and predisposing medical
25 conditions such as crescendo transient ischemic attacks, emboli and sequential strokes. Where pretreatment for

stroke or ischemia is impossible or impracticable, it is important to get the NAALADase inhibitors to the affected cells as soon as possible during or after the event. In the time period between strokes, diagnosis and treatment procedures should be minimized to save the cells from further damage and death.

COMBINATION WITH OTHER TREATMENTS

In methods of treating nervous insult (particularly acute ischemic stroke and global ischemia caused by drowning and head trauma), the NAALADase inhibitors can be co-administered with one or more therapeutic agents, preferably agents which can reduce the risk of stroke (such as aspirin), and more preferably agents which can reduce the risk of a second ischemic event (such as ticlopidine).

The NAALADase inhibitors can be co-administered with one or more therapeutic agents either (i) together in a single formulation, or (ii) separately in individual formulations designed for optimal release rates of their respective active agent. Each formulation may contain from about 0.01% to about 99.99% by weight, preferably from about 3.5% to about 60% by weight, of a NAALADase inhibitor, as well as one or more pharmaceutical excipients, such as wetting, emulsifying and pH buffering agents.

In Vivo Toxicity of NAALADase Inhibitors

To examine the toxicological effect of NAALADase inhibition in vivo, a group of mice were injected with 2-(phosphonomethyl)pentanedioic acid, a NAALADase inhibitor of high activity, in doses of 1, 5, 10, 30, 100, 300 and 500 mg/kg body weight. The mice were subsequently observed two times per day for 5 consecutive days. The survival rate at each dose level is provided below in TABLE I. The results show that the NAALADase inhibitor is non-toxic to mice, suggesting that it would be similarly non-toxic to humans when administered at therapeutically effective amounts.

TABLE I

TOXICOLOGICAL EFFECTS OF NAALADASE INHIBITORS							
Dose (mg/kg)	1	5	10	30	100	300	500
Survival Rate After 5 days (%)	100	100	100	100	100	100	66.7




In Vitro Inhibition of NAALADase Activity

Various compounds of formula I were tested for in vitro inhibition of NAALADase activity. The results are provided below in Table III.







77

TABLE II







IN VITRO INHIBITION OF NAALADASE ACTIVITY

Compound	K_i (nM)
5	
0.293 ± 0.08	
10 2-(phosphonomethyl)pentanedioic acid	
	700.00 ± 67.3
15 2-(phosphonomethyl)succinic acid	
	1.89 ± 0.19
20 2-[[[(2-carboxyethyl)hydroxyphosphinyl]-methyl]pentanedioic acid	








78

		
5		34.15
		35.85
10		54.50
		113.50
15		180.00
20		148.50
25		231.67

79

		532.00
5		1100.00
10		68.00
15		70.00
20		89.50
25		22.67

SUBSTITUTE SHEET (RULE 26)

		80	
			204.00
5			199.00
10			185.00
15			177.00
20			22.50
25			92.00
			117.00

The results show that 2-(phosphonomethyl)pentanedioic acid exhibits high NAALADase inhibiting activity, with a K_i of 0.293 nM. The activity of this compound is over 1000 times greater than that of previously described
5 NAALADase inhibitors.

By comparison, 2-(phosphonomethyl)succinic acid exhibits much lower NAALADase inhibiting activity, suggesting that a glutamate analog attached to the phosphonic acid contributes to its NAALADase inhibiting
10 activity.

The results also show that 2-[[[(2-carboxyethyl)-hydroxyphosphinyl]methyl]pentanedioic acid, which has an additional carboxylic acid side chain similar to the aspartate residue found in NAAG, exhibits a lower
15 NAALADase inhibiting activity than 2-(phosphonomethyl)-pentanedioic acid.

Protocol for Assaying In Vitro Inhibition of NAALADase
Activity

20 The amount of [3 H]Glu liberated from [3 H]NAAG in 50 mM Tris-Cl buffer was measured for 15 minutes at 37° C using 30-50 μ g of synaptosomal protein. Substrate and product were resolved by anion-exchange liquid chromatography. Duplicate assays were performed so that
25 no more than 20% of the NAAG was digested, representing the linear range of peptidase activity. Quisqualate (100








μM) was included in parallel assay tubes to confirm the specificity of the measurements.

In Vitro Assay of NAALADase Inhibitors on Ischemia

5 To examine the in vitro effect of NAALADase inhibitors on ischemia, cortical cell cultures were treated with various compounds of formula I during an ischemic insult (potassium cyanide and 2-deoxyglucose) and for one hour thereafter (for experimental details,
10 see Vornov et al., J. Neurochem, Vol. 65, No. 4, pp. 1681-1691 (1995)).


 The neuroprotective effect of each tested compound is provided below in TABLE III(a). Neuroprotective effect is expressed as EC_{50} , the concentration which is
15 required to cause a 50% reduction in glutamate toxicity following an ischemic insult.

TABLE III (a)


	Compound	EC ₅₀ (nM)
5		0.67
10		373.0
15		112.0
		132.0
20		100.0
		767.0
25		794.0

84

5




37.00




79.00

10




2.00

15




834.00

20



4315.00

25



1670.00

The dose-response of this effect, as measured by the % toxicity at different concentrations of 2-(phosphonomethyl)pentanedioic acid, is provided below in TABLE III(b) and graphically presented in FIG. 1.

5

TABLE III(b)

	<u>Dose</u>	<u>% Toxicity</u>
	Control	100.00 \pm 9.0 (n = 5)
	100 pM	66.57 \pm 4.38 (n = 5)
10	1 nM	42.31 \pm 9.34 (n = 5)
	10 nM	33.08 \pm 9.62 (n = 5)
	100 nM	30.23 \pm 9.43 (n = 5)
	1 μ M	8.56 \pm 8.22 (n = 5)

15 The results show that toxicity decreased as the concentration of 2-(phosphonomethyl)pentanedioic acid increased, suggesting that NAALADase inhibitors would be effective in treating ischemia or neuronal damage caused by ischemia.

20 The methods for this assay are described in detail below. Specifically, cell cultures were exposed to potassium cyanide and 2-deoxyglucose (2-DG) (10 mM) and analyzed for release of lactate dehydrogenase (LDH).

25

In Vitro Toxicity of NAAG

To examine the in vitro toxicity of NAAG, cortical cell cultures were treated with NAAG (in concentrations ranging from 3 μ M to 3 mM) for 20 minutes. The toxicity measurement for each concentration of NAAG is provided below in TABLE IV and graphically presented in FIG. 2.

TABLE IV

	<u>Dose of NAAG</u>	<u>% Toxicity</u>
10	3 μ M	3.51 (n = 1)
	10 μ M	4.30 \pm 3.12 (n = 3)
	30 μ M	11.40 \pm 6.17 (n = 3)
	100 μ M	12.66 \pm 5.50 (n = 3)
	300 μ M	13.50 \pm 4.0 (n = 3)
15	1 mM	21.46 \pm 4.20 (n = 3)
	3 mM	45.11 \pm 4.96 (n = 3)

The results show that toxicity increased as the concentration of NAAG increased. The toxicity is attributed to the release of glutamate by NAAG when cleaved by NAALADase.

In Vitro Assay of NAALADase Inhibitors on Toxicity of NAAG

To examine the effect of NAALADase inhibitors on in vitro toxicity of NAAG, cortical cell cultures were

treated with 2-(phosphonomethyl)pentanedioic acid (1 μ M) during exposure to NAAG and for one hour thereafter. The toxicity measurement for each concentration of NAAG is provided below in TABLE V and graphically presented in FIG. 3.

TABLE V

	<u>Dose of NAAG</u>	<u>% Toxicity</u>
10	3 μ M	-4.71 (n = 1)
	10 μ M	-3.08 \pm 0.81 (n = 3)
	30 μ M	-4.81 \pm 1.13 (n = 3)
	100 μ M	-2.87 \pm 0.78 (n = 3)
	300 μ M	-2.09 \pm 0.48 (n = 3)
15	1 mM	0.26 \pm 1.11 (n = 3)
	3 mM	16.83 \pm 8.76 (n = 3)

When compared to the results of FIG.2/TABLE IV, the results of FIG.3/TABLE V show that toxicity decreased considerably after treatment with the NAALADase inhibitor, suggesting that it would be effective in treating glutamate abnormalities.

In Vitro Assay of NAALADASE Inhibitors on Ischemia at Different Times of Administration

To examine the effect of NAALADase inhibitors on in vitro ischemic toxicity at different times of

88

administration, cortical cell cultures were treated with 2-(phosphonomethyl)pentanedioic acid (i) during an ischemic insult and for one hour thereafter (exposure and recovery); (ii) for one hour following ischemic insult (recovery only); and (iii) for one hour beginning 30 minutes after ischemic insult (delayed 30 minutes). The toxicity measurement for each time of administration is provided below in TABLE VI and graphically presented in FIG. 4.

10

TABLE VI

	<u>Time of Administration</u> <u>Relative to Ischemic Insult</u>	<u>% Toxicity</u>
	Control	100.00%
15	Exposure & Recovery	2.54%
	Recovery Only	9.03%
	Delayed 30 Minutes	31.49%

The results show that significant neuronal protection is achieved when NAALADase inhibitors are administered during exposure and recovery from an ischemic insult, and even after a 30 minute delay following the ischemic insult.

25

Protocol for In Vitro Toxicity Assaya. Cell Culture

Dissociated cortical cell cultures are prepared using the papain-dissociation method of Heuttner and
5 Baughman (1986) as modified by Murphy and Baraban (1990). See TABLE VII for the Dissociated Culture Protocol as used herein. Fetuses of embryonic day 17 are removed from timed pregnancy rats (Harlan Sprague Dawley). The cortex is rapidly dissected out in Dulbecco's phosphate-
10 buffered saline, stripped of meninges, and incubated in a papain solution for 15 minutes at 37° C. The tissue is then mechanically triturated and pelleted at 500 g (1000-2000 rpm on swinging bucket Beckman). The pellet is resuspended in a DNAase solution, triturated with a 10 ml
15 pipette x15-20, layered over a "10 x 10" solution containing albumin and trypsin inhibitor (see TABLE VII for an example of a "10 x 10" solution), repelleted, and resuspended in a plating medium containing 10% fetal bovine serum (HyClone A-1111-L), 5% heat-inactivated
20 Equine serum (HyClone A-3311-L), and 84% modified Earle's basal medium (MEM) (Gibco 51200-020) with high glucose (4.5 g/L), and 1 g/L NaHCO₃. Each 24-well plate is pretreated with poly-D-lysine (0.5 ml/well of 10 µg/ml) for 1 h and rinsed with water before plating. Cultures
25 are plated at 2.5×10^6 cells/ml with each well of a 24 well plate receiving 500 µl/well. Alternatively, 35 mm

90

dishes can be plated at 2 ml/dish, 6 well plates at 2 ml/well, or 12 well plates at 1 ml/well. After plating, 50% of the medium is changed every 3-4 days with growth serum containing 5% heat-inactivated Equine serum (HyClone A-3311-L), 95% modified Earle's basal medium (MEM) (Gibco 51200-020), and 1% L-Glutamine (Gibco 25030-081). Experiments are performed after 21 days in cultures. Cultures are maintained in a 5% CO₂ atmosphere at 37° C. These methodologies are described in further detail below in the TABLE VII.

TABLE VII

DISSOCIATED CULTURE PROTOCOL	
I. PREPARE SOLUTIONS	
Stocks/Solutions	
<p>5 <u>DNAase Stock, 1 ml</u> <u>(100x)</u> 5 mg DNAase I (Worthington LS002004); 1 ml dissoc. EBSS; 10 freeze as 50 μl aliquots.</p>	<p><u>Dulbecco's PBS, 500 ml</u> 4 gm NaCl (J.T. Baker 3624-01); 1.06 gm $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ (Fisher S373-3); 100 mg KCl (Fisher P217-500); 100 mg KH_2PO_4 (Sigma P-0662); 500 ml dH_2O; adjust pH to 7.4 and sterile filter.</p>
<p>15 <u>Dissociated EBSS, 500 ml</u> 1.1 gm NaHCO_3; 50 ml EBSS stock (Gibco 14050-025); 450 ml dH_2O; sterile filter.</p>	<p><u>EDTA Stock, 10 ml</u> 184.2 mg EDTA sodium salt (Sigma ED4S); 10 ml dH_2O; sterile filter.</p>

DISSOCIATED CULTURE PROTOCOL			
5	<table><tr><td><u>10 and 10 Stock, 10 ml</u> 100 mg BSA (Sigma A-4919); 100 mg Trypsin Inhibitor from Egg White (Sigma T-2011); 10 ml dissoc. EBSS; sterile filter.</td><td><u>Poly-D-Lysine Stock, 5 ml</u> 5 mg Poly-D-Lysine, 100-150 K (Sigma P-6407); 5 ml sterile water; keep frozen.</td></tr></table>	<u>10 and 10 Stock, 10 ml</u> 100 mg BSA (Sigma A-4919); 100 mg Trypsin Inhibitor from Egg White (Sigma T-2011); 10 ml dissoc. EBSS; sterile filter.	<u>Poly-D-Lysine Stock, 5 ml</u> 5 mg Poly-D-Lysine, 100-150 K (Sigma P-6407); 5 ml sterile water; keep frozen.
<u>10 and 10 Stock, 10 ml</u> 100 mg BSA (Sigma A-4919); 100 mg Trypsin Inhibitor from Egg White (Sigma T-2011); 10 ml dissoc. EBSS; sterile filter.	<u>Poly-D-Lysine Stock, 5 ml</u> 5 mg Poly-D-Lysine, 100-150 K (Sigma P-6407); 5 ml sterile water; keep frozen.		
10	<table><tr><td>Media</td><td></td></tr></table>	Media	
Media			

DISSOCIATED CULTURE PROTOCOL

<u>Dissociated growth, 500 ml</u>	<u>Plating media, 300 ml</u>
<p>500 ml MEM (Gibco 51200-020) containing glucose and NaHCO₃, (2.25 gm glucose and 0.5 gm NaHCO₃);</p> <p>25 ml heat-inactivated Equine Serum (HyClone A-3311-L);</p> <p>5 ml L-Glutamine (200 mM, 100x stock, Gibco 25030-081);</p> <p>sterile filter.</p>	<p>250 ml MEM containing glucose and sodium bicarbonate (2.25 gm glucose and 0.5 gm NaHCO₃ in 500 ml Gibco MEM 51200-020);</p> <p>30 ml Fetal Bovine Serum (HyClone A-1111-L).</p>
<p>15 15 ml heat-inactivated Equine Serum (HyClone A-3311-L);</p> <p>3 ml L-Glutamine (200 mM, 100x stock, Gibco 25030-081); (Gibco 15140-015);</p> <p>1 ml Penicillin-Streptomycin stock.</p>	

DISSOCIATED CULTURE PROTOCOL	
For papain dissociation:	For DNAase treatment:
4 mg Cysteine (C-8277); 25 ml dissoc. EBSS; 5 250 µl Papain stock (Worthington LS003126); place in 37°C waterbath until clear.	<u>DNAase, 5 ml</u> 4.5 ml dissoc. EBSS; 500 µl "10 and 10" stock; 50 µl DNAase stock. <u>"10 and 10", 5 ml</u> 4.5 ml of EBSS; 500 µl "10 and 10" stock.
II. COAT DISHES	
10	Use poly-d-lysine stock at 1:100 dilution to coat 24-well plates (0.5 ml/well) or at 1:10 dilution to coat 35 mm glass cover slips (1.0 ml/coverslip). Leave until end of dissection.

DISSOCIATED CULTURE PROTOCOL

III. DISSECT TISSUE

5 Use Harlan Sprague-Dawley timed pregnancy rats,
ordered to arrive at E-17.

Decapitate, spray abdomen down with 70% EtOH.

Remove uterus through midline incision and place in
sterile dPBS.

10 Remove brains from embryos, leaving them in dPBS.

15 Brain removal: Penetrate skull and skin with fine
forceps at lambda. Pull back to open posterior fossa.
Then move forceps anteriorly to separate sagittal
suture. Brain can be removed by scooping back from
olfactory bulbs under the brain.

Move brains to fresh dPBS; subsequently, dissect away
from cortex.

DISSOCIATED CULTURE PROTOCOL**IV. PAPAIN DISSOCIATION**

Transfer cortices equally to two 15 ml tubes
containing sterile papain solution, maintained at 37°
5 C.

Triturate xl with sterile 10 ml pipette.

Incubate only for 15 minutes at 37° C.

Spin at 500 G for 5 minutes (1000-2000 RPM on swinging
bucket Beckman).

10 V. DNAase TREATMENT

Remove supernatant and any DNA gel layer from cell
pellet (or pick up and remove pellet with pipette).

Move cell pellet to DNAase solution.

15 Triturate with 10 ml pipette, x15-20.

Layer cell suspension over the "10 and 10" solution by
pipetting it against the side of the tubes.

Spin again at 500 G for 5 minutes (cells will spin
into "10 and 10" layer).

20 Wash tube sides with plating media without disturbing
pellet.

Pipette off the media wash and repeat the wash.

DISSOCIATED CULTURE PROTOCOL

VI. PLATE

Add about 4.5 ml plating media to each pellet for 5 ml volume.

5 Re-suspend with 10 ml pipette.

Pool cells into a single tube.

Quickly add 10 μ l of the suspended cells to a hemocytometer so that they do not settle.

10 Count cells per large square, corresponding to 10 million cells/ml.

Put re-suspended cells into a larger container so that they number 2.5 million cells/ml.

Triturate to homogeneity.

15 Finish coating plates:

Aspirate or dump Lysine;

Wash x1 with sterile water and dump.

20 Add plating media, with cells, to the plates as follows:

35 mm dishes 2 ml/dish;

6 well plate 2 ml/well;

12 well plate 1 ml/well;

24 well plate 500 μ l/well.

DISSOCIATED CULTURE PROTOCOL**VII. FEED**

Cultures are usually made on Thursdays.

Start feeding twice a week; beginning the following Monday, feedings on Mondays and Fridays.

- 5 Remove 50% of volume and replace with fresh growth media.

b. Ischemic Insult using potassium cyanide and
10 2-deoxyglucose

Twenty-one to twenty-four days following the initial cortical cell plating, the experiment is performed. The cultures are washed three times in HEPES buffered saline solution containing no phosphate. The cultures are then
15 exposed to potassium cyanide (KCN) (5 mM) and 2-deoxyglucose (2-DG) (10 mM) for 20 minutes at 37° C. These concentrations were shown previously to induce maximal toxicity (Vornov et al., J. Neurochem, Vol. 65, No. 4, pp. 1681-1691 (1995)). At the end of 24 hours,
20 the cultures are analyzed for release of the cytosolic enzyme lactate dehydrogenase (LDH), a standard measure of cell lysis. LDH measurements are performed according to the method of Koh and Choi, J. Neuroscience Methods (1987).

c. NAAG Induced Neurotoxicity

Cultures are assessed microscopically and those with uniform neuronal densities are used in the NAAG neurotoxicity trials.

5 At the time of the experiment, the cultures are washed once in HEPES-buffered saline solution (HBSS; NaCl 143.4 mM, HEPES 5 mM, KCl 5.4 mM, MgSO₄ 1.2 mM, NaH₂PO₄ 1.2 mM, CaCl₂ 2.0 mM, D-glucose 10 mM) (Vornov et al., 1995) and then exposed to various concentrations of NAAG
10 for 20 minutes at 37° C. NAAG concentrations range from 3 μM to 3 mM, and include 3 μM, 10 μM, 30 μM, 100 μM, 300 μM, 1 mM, and 3 mM. At the end of exposure, the cells are washed once with HEPES buffered saline solution and then replaced with serum free modified Earle's basal
15 medium. The cultures are then returned to the CO₂ incubator for 24 hour recovery.

d. Lactate Dehydrogenase Assay

Release of the cytosolic enzyme lactate dehydrogenase (LDH), a standard measure of cell lysis, is
20 used to quantify injury (Koh and Choi, 1987). LDH-activity measurements are normalized to control for variability between culture preparations (Koh and Choi, 1987). Each independent experiment contains a control condition in which no NAALADase inhibitors are added; a
25 small amount of LDH activity is found in these controls. This control measurement is subtracted from each

100

experimental point. These values are normalized within each experiment as a percentage of the injury caused by NAAG/ischemia. Only main effects of NAALADase inhibitors are considered; interactions between dose and condition are not examined statistically.

A measurement of the potency of each compound tested is made by measuring the percentage of LDH release into the growth media after exposure to NAAG/ischemia plus NAALADase inhibitor or NAAG/ischemia plus saline (control). Since high concentrations of glutamate may be toxic to cells in certain circumstances, measurement of glutamate toxicity is observed using LDH as a standard measurement technique.

In Vivo Assay of NAALADase Inhibitors on Cortical Injury following MCAO in SHRSP Rats

To examine the effect of NAALADase inhibitors on cortical injury in vivo, the infarct volume was measured in SHRSP rats which had sustained middle cerebral artery occlusion (MCAO) and had subsequently been treated with (i) saline; (ii) 10 mg/kg of 2-(phosphonomethyl)-pentanedioic acid followed by 2 mg/kg/hr of 2-(phosphonomethyl)pentanedioic acid for 1 hour; or (iii) 100 mg/kg of 2-(phosphonomethyl)pentanedioic acid followed by 20 mg/kg/hr of 2-(phosphonomethyl)-pentanedioic acid for one hour.

The cortical injury volume for each group of rats is provided below in TABLE VIII and graphically presented in FIG. 5.

5

TABLE VIII

10

15

Cortical Injury Volume (mm ³)±S.E.M.	
Control	184.62±33.52 (n = 10)
10 mg/kg	135.00±32.18 (n = 10)
100 mg/kg	65.23±32.18 (n = 10)
Cortical Injury Volume (% injury)±S.E.M.	
Control	34.44±6.53 (n = 10)
10 mg/kg ³	29.14±7.68 (n = 10)
100 mg/kg	13.98±6.64 (n = 10)
Cortical Protection (% protection)	
Control	0%
10 mg/kg	27%
100 mg/kg	65%

The results show that cortical injury volume decreased and cortical protection increased as the amount of NAALADase inhibitor increased, further supporting the neuroprotective effect of the NAALADase inhibitor.

Protocol for In Vivo Assay of NAALADase Inhibitors on
Cortical Injury

A colony of SHRSP rats is bred at Johns Hopkins School of Medicine from three pairs of male and female rats obtained from the National Institutes of Health (Laboratory, Sciences Section, Veterinary Resources Program, National Center for Research Resources, Bethesda, MD). All rats are kept in a virus-free environment and maintained on regular diet (NIH 31, Zeigler Bros, Inc.) with water ad libitum. All groups of rats are allowed to eat and drink water until the morning of the experiment.

Transient occlusion of the middle cerebral artery (MCA) is induced by advancing a 4-0 surgical nylon suture into the internal carotid artery (ICA) to block the origin of the MCA (Koizumi, 1986; Longa, 1989; Chen, 1992). The rats are anesthetized with 4% halothane, and maintained with 1.0% to 1.5% halothane in air enriched oxygen using a face mask. Rectal temperature is maintained at $37.0 \pm 0.5^{\circ}\text{C}$ throughout the surgical procedure using a heating lamp. The right femoral artery is cannulated for measuring blood gases (pH, oxygen tension $[\text{PO}_2]$, carbon dioxide tension $[\text{PCO}_2]$) before and during ischemia, for monitoring blood pressure during the surgery. The right common carotid artery (CCA) is exposed through a midline incision; a self-retraining

retractor is positioned between the digastric and mastoid muscles, and the omohyoid muscle is divided. The right external carotid artery (ECA) is dissected and ligated. The occipital artery branch of the ECA is then isolated
5 and coagulated. Next, the right internal carotid artery (ICA) is isolated until the pterygopalatine artery is exposed, and carefully separated from the adjacent vagus nerve. The pterygopalatine artery is ligated with 4-0 silk suture close to its origin.

10 After the CCA is ligated with 4-0 silk suture, a 4-0 silk suture to prevent bleeding from a puncture site, through which a 2.5 cm length of 4-0 monofilament nylon suture (Ethilon), its tip rounded by heating near a electric cautery, is introduced into the ICA lumen. A 6-
15 0 silk suture is tightened around the intraluminal nylon suture at the bifurcation to prevent bleeding, and the stretched sutures at the CCA and the ICA are released. The nylon suture is then gently advanced as far as 20 mm.

Anesthesia is terminated after 10 minutes of MCA
20 occlusion in both groups, and the rats were awakened 5 minutes thereafter. After 2 hours of ischemia, anesthesia is reanesthetized, and reperfusion is performed by withdrawing the intraluminal nylon suture until the distal tip became visible at the origin of the
25 ICA.

Arterial pH and PaCO_2 , and partial pressure of

oxygen (PaO_2) are measured with a self-calibrating Radiometer electrode system (ABL 3; Copenhagen, Denmark). Hemoglobin and arterial oxygen content are measured with a hemoximeter (Radiometer, Model OSM3; Copenhagen, Denmark). Blood glucose is measured with a glucose analyzer (model 2300A, Yellow Springs Instruments, Yellow Springs, OH).

Each group is exposed to 2 hours of right MCA occlusion and 22 hours of reperfusion. All variables but the rectal temperature are measured at baseline, at 15 minutes and 45 minutes of right MCA occlusion. The rectal temperature is measured at baseline, at 0 and 15 min of MCA occlusion, and at 0 and 22 hours of reperfusion.

In Vivo Assay of NAALADase Inhibitors on Brain Injury following MCAO in Sprague-Dawley Rats

To examine the neuroprotective effect of NAALADase inhibitors on brain injury in vivo, Sprague-Dawley rats were treated with a vehicle or 2-(phosphonomethyl)pentanedioic acid before and after sustaining a 2 hour transient middle cerebral artery occlusion (MCAO). In the control group ($n = 8$), the rats received an IP injection of saline 30 minutes post-occlusion followed by IV saline infusion at a rate of 0.5 ml/hr. In the drug treated groups, the rats received an

IP injection of 2-(phosphonomethyl)pentane-dioic acid at a dose of 100 mg/kg at 20 minutes pre-occlusion (n = 5), 30 minutes post-occlusion (n = 9), 60 minutes post-occlusion (n = 7), or 120 minutes post-occlusion (n = 4), followed by a 20 mg/kg/hr IV infusion for 4 hours (infusion rate = 0.5 ml/hr). There was a 15 minute delay between IP and IV treatments for each rat. Twenty two hours following the reperfusion, the rats were euthanized and their brains were removed. Seven coronal sections (2 mm thick) were taken and stained with 1% solution of 2,3,5-triphenyltetrazolium chloride (TTC) for 20 minutes and then fixed in 10% formalin. The anterior and posterior surface of the most rostral brain section and the posterior surface of each of the other 6 sections were imaged. The quantification of infarct size of each brain was obtained using a computer aided-digital imaging analysis system (LOATS). The brain regions completely lacking TTC-staining were characterized as representative of infarcted tissue. The total infarct volume for each rat was calculated by numeric integration of the respective sequential brain areas.

The total infarct volume for each group of rats is graphically presented in FIG. 6.

Vehicle treated rats exhibited a mean total brain infarct volume of $293 \pm 26 \text{ mm}^3$. Rats treated with 2-(phosphonomethyl)pentanedioic acid either before or after

the ischemic insult exhibited significantly lower mean total brain infarct volumes of $122 \pm 26 \text{ mm}^3$ ($p = 0.003$ vs. vehicle) for 20 minute pre-treatment, $208 \pm 40 \text{ mm}^3$ ($p = 0.2$ vs. vehicle) for 30 minute post-treatment, 125 ± 57 5 mm^3 ($p = 0.015$ vs. vehicle) for 60 minute post-treatment, and $133 \pm 35 \text{ mm}^3$ ($p = 0.005$ vs. vehicle) for 120 minute post-treatment. These results indicate that 2-(phosphonomethyl)pentanedioic acid is neuroprotective in rat MCAO model of stroke when administered up to 2 hours 10 post-occlusion.

Protocol for In Vivo Assay of NAALADase Inhibitors on
Brain Injury

Male Sprague-Dawley rats (260-320 g) were used. 15 Prior to the experiment, the rats were individually housed and allowed free access to food and water. Each rat received two surgeries: jugular vein cannulation for IV infusion and MCAO. During surgeries, the rat was anesthetized with 2% halothane delivered in oxygen via an 20 inhalation mask. The body temperature was monitored and regulated at normothermic level using a homeothermic heating system. First, a PE-50 polyethylene catheter was inserted into the right jugular vein. One hour later, the rat was reanesthetized for MCAO surgery. The MCAO 25 was achieved using the endovascular suture method described by Long et al., Stroke, Vol. 20, pp. 84-91

(1989). Specifically, the right external carotid artery (ECA) was exposed, coagulated and transected. A 3-0 monofilament nylon suture with a blunted tip was introduced into the proximal stump of the ECA via an arteriotomy and advanced 20 mm from the carotid bifurcation until it lodged in the proximal region of the anterior cerebral artery, thereby occluding the origin of the MCA. The rats were allowed to wake up; 2 hours later, the rats were reanesthetized for reperfusion, during which the nylon suture was retracted to the stump of the ECA allowing blood recirculation to the MCA.

In Vivo Assay of NAALADase Inhibitors on Stroke-Induced Rise in Brain Glutamate Levels

To examine the effect of NAALADase inhibitors on hyperglutamatergic disorders in vivo, rats with stroke-induced rise in brain glutamate levels were treated with a vehicle or 2-(phosphonomethyl)pentanedioic acid.

The results are graphically presented in FIGS. 7, 8 and 9.

The results show that 2-(phosphonomethyl)-pentanedioic acid treatment (100 mg/kg IP followed by 20 mg/kg/hr IV) significantly attenuated stroke-induced extracellular glutamate increases in the striatum (FIG. 7) as compared to vehicle treated rats ($p < 0.05$), and completely prevented concurrent glutamate changes in the

parietal cortex ($p < 0.01$; FIG. 8). In contrast, there was no significant effect of the stroke itself on glutamate in the frontal cortex and no subsequent difference between the vehicle and 2-(phosphonomethyl)-
5 pentanedioic acid treated groups (FIG. 9). Values are expressed as % baseline where baseline constitutes the mean of three consecutive 20 minute samples preceding stroke. Absolute basal (pretreatment) values for glutamate (mean \pm SEM) in caudate, parietal and frontal
10 cortices were 0.25 ± 0.1 , 1.1 ± 0.3 and 0.6 ± 0.1 μM , respectively, in the vehicle treated rats, and 0.46 ± 0.1 , 2.0 ± 0.7 and 0.9 ± 0.3 μM , respectively, in the 2-(phosphonomethyl)pentanedioic acid treated rats.

15 Protocol for In Vivo Assay of NAALADase Inhibitors on
Stroke-Induced Rise in Brain Glutamate Levels

Male Sprague Dawley rats (270-330 g, $n = 5-6$ per group) were implanted with concentric microdialysis probes similar to previously described procedures
20 (Britton et al., J. Neurochem., Vol. 67, pp. 324-329 (1996)). In brief, under halothane anaesthesia, probes (constructed in-house using Cuprophane capillary membrane; 10K mw cut off; 2 mm dialyzing length) were implanted into the frontal cortex (AP = +3.5; ML = 3; DIV
25 = 3), caudate nucleus (AP = 0; ML = 3; DV = 6.6), and parietal cortex (AP = -2; ML = 5; DV = 3) (coordinates in

mm relative to bregma and dura, respectively), regions believed to represent core and penumbral areas of ischemia-induced injury. Glutamate levels in dialysate were determined using precolumn o-phthaldialdehyde derivatization, followed by HPLC with fluorometric detection.

Approximately 20 hours after probe implantation, the rats were dialyzed with perfusion fluid (125 mM NaCl, 2.5 mM KCl, 1.18 mM MgCl₂, and 1.26 mM CaCl₂) at a rate of 2.5 μ l/min. Following a 60 minute stabilization period, dialysis samples were collected every 20 minutes. After collecting 3 baseline samples, the rats were anaesthetized with halothane and subjected to temporary ischemia using the filament method of MCAO (Britton et al., Life Sciences, Vol. 60, No. 20, pp. 1729-1740 (1997)). In brief, the right external carotid artery (ECA) was exposed and its branches coagulated. A 3-0 monofilament nylon suture was introduced into the internal carotid artery via an arteriotomy in the ECA and advanced until it lodged in the proximal region of the anterior cerebral artery, thus occluding the origin of the MCA. The endovascular suture was retracted to allow reperfusion 2 hours after occlusion.

Body temperature was maintained normothermic throughout stroke surgery and reperfusion procedures. The rats were dosed IP with 100 mg/kg 2-

110

(phosphonomethyl)pentanedioic acid at -20 minute pre-occlusion and IV with 20 mg/kg/hr for 4 hours at the time of occlusion. Dialysis samples were collected every 20 minutes from unanesthetized rats. Following 24 hours of
5 reperfusion, the rats were sacrificed, their brains were removed, and 7 coronal sections (2 mm thick) were taken from the region beginning 1 mm from the frontal pole and ending just rostral to the cortico-cerebellar junction. Analysis of ischemic cerebral damage was achieved using
10 TTC staining and computer assisted image analysis as described by Britton et al. (1997), supra.

In Vivo Assay of NAALADase Inhibitors on Myelin
Formation Following Sciatic Nerve Cryolesion

15 It was recently demonstrated that NAALADase is down-regulated in glial cells as they start to form myelin and is absent in myelinating Schwann cells. Based on this data, the inventors hypothesized that inhibition of NAALADase may affect the signaling mechanism between
20 axons and Schwann cells and result in increasing myelination. To test this hypothesis, the inventors examined the effect of 2-(phosphonomethyl)pentanedioic acid on nerve regeneration and myelination following cryolesion of the sciatic nerve in male mice.

25 The results are provided below in TABLE IX and graphically presented in FIG. 10(a) and FIG. 10(b).

111

TABLE IX

IN VIVO EFFECT OF NAALADASE INHIBITORS ON MYELIN
FORMATION FOLLOWING SCIATIC NERVE CRYOLESION

5	2- (phosphonomethyl) - pentanedioic acid	vehicle
10	ratio of # of myelinated axons (drug/vehicle)	1.5
15	# of myelinated lamellae (ave. + SEM)	16.53 \pm 0.65
	% increase of myelinated lamellae over vehicle	13.77 \pm 0.09
20	significance by t-test	p < 0.005

As detailed in FIG. 10(a) and FIG. 10(b), both light
and transmission electron microscopy (TEM) examination of
the nerve 3 mm distal to the site of cryolesion

112

demonstrated a significant increase in the number of myelinated axons (1.5-fold increase) and myelin thickness (20% increase, $p < 0.005$), as compared to nerves in mice treated with vehicle.

5 FIG. 10(a) and FIG. 10(b) show a photomicrograph of this effect. Sections were stained with toluidine blue which stains myelin. Sciatic nerves treated with 2-(phosphonomethyl)-pentanedioic acid containing implants, compared with sciatic nerves treated with vehicle
10 containing implants, exhibited an increase in myelinated axon number as well as an increase in myelin thickness.

Protocol for In Vivo Assay of NAALADase Inhibitors on
Myelin Formation Following Sciatic Nerve Cryolesion

15 Cryolesion of the mouse sciatic nerve was performed according to Koenig et al., Science, Vol. 268, pp. 1500-1503 (June 1995). In brief, each mouse was anesthetized and its sciatic nerve was exposed in the upper thigh and cryolesioned using a copper cryode (diameter = 0.5 mm)
20 that was dipped in liquid nitrogen and repeatedly applied to the upper part of the nerve. The extent of the lesion was approximately 1 mm.

2- (Phosphonomethyl)pentanedioic acid was incorporated into silicone strips according to the method
25 of Connold et al., Developmental Brain Res, Vol. 28, pp. 99-104 (1986), and was implanted at the site of

113

cryolesion on day 0 and replaced on days 3, 6, 9 and 12. Approximately 2.5 μ g/day of 2-(phosphonomethyl)-pentanedioic acid was released from the silicone implants each day. Both right and left sciatic nerves of each mouse were lesioned; right-side nerves were treated with silicone implant strips containing vehicle alone while left-side nerves were treated with silicone implants containing 2-(phosphonomethyl)pentanedioic acid. Fifteen days after surgery, the mice were sacrificed and their sciatic nerve segments were collected and processed for light microscopy and TEM analysis. Randomly chosen fields 2-3 mm distal to the lesion were qualitatively analyzed by light microscopy using 1-micrometer-thick toluidine blue stained cross sections and photographic images were captured.

In Vivo Assay of NAALADase Inhibitors on Parkinson's Disease

To examine the effect of NAALADase inhibitors on Parkinson's Disease in vivo, MPTP lesioned mice were treated with 2-(phosphonomethyl)pentanedioic acid or a vehicle.

The percent of dopaminergic neurons for each group of mice is provided below in TABLE X and graphically presented in FIG. 11.

114

TABLE X

IN VIVO EFFECT OF NAALADASE INHIBITORS ON PARKINSON'S
DISEASE

5		Percent Strial TH
		Innervation Density (mean \pm SEM)
10	vehicle/vehicle	24.74 \pm 1.03
	MPTP/vehicle	7.82 \pm 0.68
	MPTP/2-(phosphonomethyl)- pentanedioic acid	16.28 \pm 0.98
15		

Mice treated with MPTP and vehicle exhibited a substantial loss of functional dopaminergic terminals as compared to non-lesioned mice (approximately 68% loss). Lesioned mice receiving 2-(phosphonomethyl)pentanedioic acid (10 mg/kg) showed a significant recovery of TH-stained dopaminergic neurons ($p < 0.001$). These results indicate that 2-(phosphonomethyl)pentanedioic acid protects against MPTP-toxicity in mice.

25

Protocol for In Vivo Assay of NAALADase Inhibitors on
Parkinson's Disease

MPTP lesioning of dopaminergic neurons in mice was used as an animal model of Parkinson's Disease, as described by Steiner, Proc. Natl. Acad. Sci., Vol. 94, pp. 2019-2024 (March 1997). In brief, four week old male CD1 white mice were dosed IP with 30 mg/kg of MPTP for 5 days. 2-(Phosphonomethyl)pentanedioic acid (10 mg/kg) or a vehicle was administered SC along with the MPTP for 5 days, as well as for an additional 5 days following cessation of MPTP treatment. At 18 days following MPTP treatment, the mice were sacrificed and their brains were removed and sectioned. Immunostaining was performed on sagittal and coronal brain sections using anti-tyrosine hydroxylase (TH) antibodies to quantitate survival and recovery of dopaminergic neurons.

In Vivo Assay of NAALADase Inhibitors on Dynorphin-
Induced Spinal Cord Injury

To examine the neuroprotective effect of NAALADase inhibitors on excitotoxic spinal cord injury in vivo, rats which had sustained dynorphin-induced spinal cord injury were treated with a vehicle or 2-(phosphonomethyl)pentanedioic acid.

The results are graphically presented in FIG. 12. When co-administered with dynorphin A, 2-

116

(phosphonomethyl)pentanedioic acid (4 μ moles) caused significant improvement in motor scores by 24-hour post-injection, as compared to vehicle treated rats ($p < 0.05$, Kruskal-Wallis comparison). The rats were characterized
5 as ambulatory or not on the basis of their assigned neurological scores (0 to 4). At 24 hours post-injection, 73% of the 15 rats co-treated with 2-(phosphonomethyl)pentanedioic acid were ambulatory, in contrast to 14% of the 14 vehicle co-treated rats ($p <$
10 0.05). These results indicate that 2-(phosphonomethyl)-pentanedioic acid provides effective protection against dynorphin-induced spinal cord injury.

Protocol for In Vivo Assay of NAALADase Inhibitors on
15 Dynorphin-Induced Spinal Cord Injury
Spinal Subarachnoid Injections

Dynorphin-induced spinal cord injury was performed according to Long et al., JPET, Vol. 269, No. 1, pp. 358-366 (1993). In brief, spinal subarachnoid injections
20 were delivered using 30-gauge needles inserted between the L4-L5 vertebrae of male Sprague-Dawley rats (300-350 g). The rats were anesthetized with halothane and dorsal midline incisions were made immediately rostral to the pelvic girdle. By using the vertebral processes as
25 guides, the needle was advanced to pass into the subarachnoid space surrounding the cauda equina. Correct

needle placement was verified by CSF flow from the needle after its insertion. Injections were delivered using a Hamilton microsyringe in a total volume of 20 μ l which contained dynorphin (20 nmol), the cannula flush and 2-
5 (phosphonomethyl)pentanedioic acid or vehicle. After injections, the incisions were treated with the topical antibacterial furazolidone and closed with wound clips. Rapid recovery from the halothane anesthesia enabled neurological evaluations to be made within 5 minutes of
10 injections.

Neurological Evaluations

Neurological function was evaluated using a 5-point ordinal scale, with scores being assigned as follows: 4 = normal motor function; 3 = mild paraparesis, with the
15 ability to support weight and walk with impairment; 2 = paraparesis, with the ability to make walking movements without fully supporting weight; 1 = severe paraparesis, in which rats could make limited hind limb movement, but not walking movement; and 0 = flaccid paralysis, with
20 complete absence of any hind limb movement. Neurological evaluations were made 24 hours after dynorphin A injection.

Statistics

Differences in the neurological scores among
25 treatment groups were determined by means of the Mann-Whitney U test or the Kruskal-Wallis test.

In Vitro Assay of NAALADase Inhibitors on Amyotrophic
Lateral Sclerosis (ALS)

To examine the neuroprotective effect of NAALADase inhibitors on Amyotrophic Lateral Sclerosis (ALS), spinal
5 cord organotypic cultures were treated with
threohydroxyaspartate (THA), 2-(phosphonomethyl)-
pentanedioic acid, or THA combined with 2-
(phosphonomethyl)pentanedioic acid, and assayed for
choline acetyltransferase (ChAT) activity.

10 The ChAT activity for each treatment of the spinal
cord organotypic cultures is provided below in TABLE XI
and graphically presented in FIG. 13.

119

TABLE XI

NEUROPROTECTIVE EFFECT OF NAALADASE INHIBITORS IN
SPINAL CORD CULTURE MODEL OF ALS

5	Treatment	ChAT Activity (% of Control)
	control	100 \pm 22.1
10	2-(phosphonomethyl)- pentanedioic acid alone	108 \pm 18.4
	THA alone	36 \pm 12.1
15	2-(phosphonomethyl)- pentanedioic acid and THA	121 \pm 18.8
20		

As shown in FIG. 13, treatment of the spinal cord organotypic cultures with 100 μ M THA resulted in a reduction of ChAT activity to approximately 36% of control cultures. Co-incubation of the cultures with THA and 2-(phosphonomethyl)pentanedioic acid (100 nM - 10 μ M) significantly protected the cultures from THA toxicity.

120

The dose-response of this effect is provided below in TABLE XII and graphically presented in FIG. 14.

TABLE XII
5 NEUROPROTECTIVE EFFECT OF NAALADASE INHIBITORS IN
SPINAL CORD CULTURE MODEL OF ALS

	ChAT Activity (% of Control)
10 control	100.0
THA	0
15 THA and 1 nM 2-(phosphonomethyl)- pentanedioic acid	-23.9 ± 18.6
THA and 10 nM 2-(phosphonomethyl)- pentanedioic acid	23.1 ± 12.5
20 THA and 100 nM 2-(phosphonomethyl)- pentanedioic acid	87.5 ± 21.7
25 THA and 1 μM 2-(phosphonomethyl)- pentanedioic acid	187.7 ± 32.8

121

THA and 10 μ M 2-(phosphonomethyl)-pentanedioic acid 128.7 \pm 17.2

5 Spinal cord cultures were incubated with various doses of 2-(phosphonomethyl)pentanedioic acid (1 nM to 10 μ M) in the presence of THA (100 μ M) for 14 days. As shown in FIG. 14, 2-(phosphonomethyl)pentanedioic acid exhibited dose-dependent protection against THA-induced
10 toxicity with maximal effects at 1 μ M.

Protocol for In Vivo Assay of NAALADase Inhibitors on
Amyotrophic Lateral Sclerosis (ALS)

Spinal Cord Organotypic Cultures

15 Organotypic cultures were prepared from lumbar spinal cord of 8 day old rats, as described by Rothstein et al., J. Neurochem., Vol. 65, No. 2 (1995), and Rothstein et al., Proc. Natl. Acad. Sci. USA, Vol. 90, pp. 6591-6595 (July 1993). In brief, lumbar spinal cords
20 were removed and sliced into 300 μ M-thick-dorsal-ventral sections, and five slices were placed on Millipore CM semipermeable 30-mm-diameter membrane inserts. The inserts were placed on 1 ml of culture medium in 35-mm-diameter culture wells. Culture medium consisted of 50%
25 minimal essential medium and phosphate-free HEPES (25 mM), 25% heat-inactivated horse serum, and 25% Hanks'

122

balanced salt solution (GIBCO) supplemented with D-glucose (25.6 mg/ml) and glutamine (2 mM), at a final pH of 7.2. Antibiotic and antifungal agents were not used. Cultures were incubated at 37° C in 5% CO₂ containing
5 humidified environment (Forma Scientific). Culture medium, along with any added pharmacological agents, was changed twice weekly.

Chronic Toxicity Model with THA

For all experiments, cultures were used 8 days after
10 preparation at which time threohydroxyaspartate (THA; 100 μ M), 2-(phosphonomethyl)pentanedioic acid (100 pM - 10 μ M), or THA (100 μ M) \pm 2-(phosphonomethyl)pentanedioic acid (100 pM - 10 μ M) were added to the culture medium. Drugs were incubated for an additional 13 to 20 days with
15 the 100 μ M THA. At the end of this period, cultures were collected assayed for ChAT activity as described below.

ChAT Assays

To determine choline acetyltransferase (ChAT) activity, the spinal cord tissues in each dish (five
20 slices) were pooled and frozen (-75° C) until assay. ChAT activity was measured radiometrically by described methods using [³H]acetyl-CoA (Amersham; Fonnum, 1975). Protein content of tissue homogenate was determined by a Coomassie Protein Assay kit (Pierce, Rockford, IL).

25

123

EXAMPLES

The following examples are illustrative of the present invention and are not intended to be limitations thereon. Unless otherwise indicated, all percentages are based upon 100% by weight of the final composition.

EXAMPLE 1

Preparation of 2-[(methylhydroxyphosphinyl)methyl]
pentanedioic acid

10 Scheme IV: $R = CH_3$, $R_1 = CH_2Ph$

Methyl-O-benzylphosphinic acid

Dichloromethylphosphite (10.0 g, 77 mmol) in 80 mL of dry diethyl ether was cooled to $-20^{\circ}C$ under an atmosphere of nitrogen. A solution of benzyl alcohol (23 g, 213 mmol) and triethylamine (10.2 g, 100 mmol) in 40 mL of diethyl ether was added dropwise over 1 hour while maintaining an internal temperature range of $0^{\circ}C$ to $10^{\circ}C$. Once addition was complete the mixture was warmed to room temperature and stirred overnight. The mixture was filtered and the solid cake washed with 200 mL of diethyl ether. The organics were combined and evaporated under reduced pressure to give 25 g of a clear and colorless liquid. The liquid was purified by flash chromatography and eluted with a 1:1 hexane/ethyl acetate to ethyl acetate gradient. The desired fractions were collected

124

and evaporated to give methyl-Q-benzylphosphinic acid (1, R = CH₃, R₁ = CH₂Ph, 6.5 g, 50%) as a clear and colorless oil. R_f 0.1 (1:1, Hexane/EtOAc).

¹H NMR (d6-DMSO): 7.4 ppm (m, 5H), 7.1 ppm (d, 1H), 5.0
5 ppm (dd, 2H), 1.5 ppm (d, 3H)

2,4-Di(benzyloxycarbonyl)butyl(methyl)-Q-benzylphosphinic acid

Methyl-Q-benzylphosphinic acid (3.53 g, 20.7 mmol)
10 in 200 mL of dichloromethane was cooled to -5° C under an atmosphere of nitrogen. Triethylamine (3.2 g, 32 mmol) was added via syringe followed by trimethylsilyl chloride (2.9 g, 27 mmol). The reaction mixture was stirred and warmed to room temperature over 1 hour. Dibenzyl 2-
15 methylenepentanedioate (2, 6.0 g, 18.5 mmol) in 10 mL of dichloromethane was added. The mixture was then stirred at room temperature overnight. The reaction mixture was cooled to 0° C and trimethylaluminum (9 mL, 18 mmol, 2.0 M in dichloromethane) was added. The flask was warmed
20 and stirred for 72 hours. The clear light yellow solution was cooled to 5° C and quenched by the slow addition of 5% hydrochloric acid. The quenched reaction mixture was warmed to room temperature and the organic layer removed. The organic layer was washed with 5%
25 hydrochloric acid and with water. The organics were dried (MgSO₄) and evaporated under reduced pressure to

125

give 8 g of a clear light yellow oil. The oil was purified on silica gel and eluted with a gradient of 1:1 hexanes/ethyl acetate to 100% ethyl acetate. The desired fractions were collected and evaporated to give 2,4-
5 di(benzyloxycarbonyl)butyl(methyl)-O-benzylphosphinic acid (3, R = CH₃, R₁ = CH₂Ph, 0.8 g, 8%) as a clear and colorless oil. R_f 0.5 (ethyl acetate).

¹H NMR (CDCl₃): 7.4 ppm (m, 15H), 5.1 ppm (m, 6H), 3.0 ppm (m, 1H), 2.4 ppm (m, 3H), 2.1 ppm (m, 3H), 1.5 ppm
10 (dd, 3H).

Elemental Analysis

Calculated C₂₈H₃₁O₆P, 0.5 H₂O: C, 68.01; H, 6.32.

Found: C, 66.85; H, 6.35.

15 2-[(Methylhydroxyphosphinyl)methyl]pentanedioic acid

2,4-di(benzyloxycarbonyl)butyl(methyl)-O-benzylphosphinic acid (0.8 g, 1.6 mmol) in 20 mL of water containing 100 mg of 10% Pd/C was hydrogenated at 40 psi for 4 hours. The mixture was filtered over a pad of
20 Celite and evaporated at high vacuum to give 2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid (4, R = CH₃, 0.28 g), 78% as a clear and colorless viscous oil.

¹H NMR (D₂O): 2.5 ppm (m, 1H), 2.2 ppm (t, 2H), 2.0 ppm (m, 1H), 1.7 ppm (m, 3H), 1.3 ppm (d, 3H).

25 Elemental Analysis

Calculated C₇H₁₃O₆P, 0.2 H₂O: C, 36.92; H, 5.93.

126

Found: C, 37.06; H, 6.31.

EXAMPLE 2**Preparation of 2-[(butylhydroxyphosphinyl)methyl]**

5

pentanedioic acid

Scheme IV: R = n-butyl, R₁ = H

Butylphosphinic Acid

Diethyl chlorophosphite (25 g, 0.16 mol) in 60 mL of dry
10 ether was cooled to 0° C under an atmosphere of nitrogen.
Butylmagnesium chloride (80 mL, 0.16 mol, 2.0 M solution
in ether) was added dropwise over a period of 2 hours
while maintaining the internal temperature at 0° C. Once
addition was complete the thick white slurry was heated
15 to 30° C for 1 hour. The suspension was filtered under
a nitrogen atmosphere and the filtrate evaporated under
reduced pressure. The clear light yellow liquid was then
brought up in 15 mL of water and stirred at room
temperature. Concentrated hydrochloric acid (0.5 mL) was
20 then added and an exothermic reaction was observed. The
mixture was stirred an additional 15 minutes and
extracted with two 75 mL portions of ethyl acetate. The
organics were combined, dried (MgSO₄) and evaporated to
give a clear and colorless liquid. The liquid was
25 treated with NaOH (40 mL, 2.0 M) and stirred for 1 hour.
The mixture was then washed with diethyl ether and

127

acidified to pH 1.0. The desired material was extracted from the acidified extract with two 100 mL portions of ethyl acetate. The organics were combined, dried (MgSO_4) and evaporated under reduced pressure to give
5 butylphosphinic acid (1, R = n-butyl, R_1 = H, 10 g, 51%) as a clear and colorless liquid.

^1H NMR (d_6 -DMSO): 6.9 ppm (d, 1H), 1.6 ppm (m, 2H), 1.4 ppm (m, 4H), 0.9 ppm (t, 3H).

10 Butyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid

Butylphosphinic acid (2.0 g, 16 mmol) in 80 mL of dry dichloromethane was cooled to 0° C under an atmosphere of nitrogen. Triethylamine (6.7 g, 66 mmol) was added followed by trimethylsilyl chloride (58 mL, 58
15 mmol, 1.0 M in dichloromethane). The mixture was stirred at 0° C for 10 minutes and dibenzyl 2-methylenepentanedioate (2) (6.4 g, 20 mmol) in 20 mL of dichloromethane was added. The cold bath was removed and the reaction warmed to room temperature and stirred
20 overnight. The mixture was then cooled to 0° C and quenched by the slow addition of 5% hydrochloric acid (50 mL). The dichloromethane layer was then removed and washed with 5% hydrochloric acid and with brine. The organic layer was dried (MgSO_4) and evaporated to give a
25 clear light golden liquid. The liquid was purified by flash chromatography and eluted with 3:1 hexane/ethyl

128

acetate containing 5% acetic acid. The desired fractions were combined and evaporated to give butyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid (3, R = n-butyl, R₁ = H) (2.9 g, 40%) as a clear and colorless oil.

5 R_f 0.12 (3:1 Hexane/EtOAc, 5% AcOH).

¹H NMR (d₆-DMSO): 7.3 ppm (m, 10H), 5.0 ppm (s, 4H), 2.7 ppm (m, 1H), 2.3 ppm (t, 2H), 1.8 ppm (m, 2H), 1.3 ppm (m, 4H), 0.8 ppm (t, 3H).

10 2-[(Butylhydroxyphosphinyl)methyl]pentanedioic acid

Butyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid (2.9 g, 6.5 mmol) in 30 mL of water containing 0.32 g 10% Pd/C was hydrogenated at 40 psi for 4.5 hours. The mixture was filtered through a pad of Celite and
15 evaporated under high vacuum to give 2-[(butylhydroxyphosphinyl)methyl]-pentanedioic acid (4, R = n-butyl) (0.75 g, 43%) as a clear and colorless viscous oil.

¹H NMR (D₂O): 2.4 ppm (m, 1H), 2.1 ppm (t, 2H), 1.9 ppm
20 (m, 1H), 1.6 ppm (m, 3H), 1.4 ppm (m, 2H), 1.1 ppm (m, 4H), 0.6 ppm (t, 3H).

Elemental Analysis

Calculated C₁₀H₁₉O₆P, 0.5 H₂O: C, 43.64; H, 7.32.

Found: C, 43.25; H, 7.12.

25

EXAMPLE 3

Preparation of 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid

Scheme IV: $R = CH_2Ph$, $R_1 = H$

5

Benzylphosphinic acid

Diethylchlorophosphite (25 g, 0.16 mol) in 100 mL of dry diethyl ether was cooled to 0° C under an atmosphere of nitrogen. Benzylmagnesium chloride (80 mL, 0.16 mol, 2.0 M solution in Et₂O) was added dropwise over two hours while maintaining a temperature below 10° C. A thick white slurry formed and stirring was continued at room temperature for 1 hour. The mixture was filtered under a nitrogen atmosphere and the filtrate evaporated under reduced pressure to give a clear and colorless liquid. The liquid was stirred as 15 mL of water was added followed by 0.5 ml concentrated hydrochloric acid. An exothermic reaction was observed and stirring was continued for an additional 30 minutes followed by extraction with ethyl acetate. The organics were combined, washed with brine, dried (MgSO₄) and evaporated. The clear light golden liquid was added to sodium hydroxide (50 mL, 2.0 M NaOH), stirred for 1 hour and washed with diethyl ether. The aqueous layer was acidified to pH 1.0 with concentrated hydrochloric acid and extracted with ethyl acetate. The organics were

130

combined, dried (MgSO_4) and evaporated to give benzylphosphinic acid (1, $\text{R} = \text{CH}_2\text{Ph}$, $\text{R}_1 = \text{H}$) (8 g, 32%) as a clear light golden oil.

^1H NMR (d_6 -DMSO): 7.3 ppm (m, 5H), 6.9 ppm (d, 1H), 3.15 ppm (d, 2H).

Benzyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid

Benzylphosphinic acid (2.3 g, 15 mmol) in 150 mL of dry dichloromethane was cooled to 0°C under a nitrogen atmosphere. Triethylamine (6.5 g, 65 mmol) was added followed by trimethylsilyl chloride (5.8 g, 54 mmol) while the reaction temperature was maintained at 0°C . After 30 minutes dibenzyl 2-methylene-pentanedioate (2) (4.4 g, 13.6 mmol) in 20 mL of dichloromethane was added over 5 minutes. The reaction mixture was allowed to warm to room temperature and stirred overnight. The clear solution was cooled to 0°C and quenched with 5% hydrochloric acid followed by removal of the organic layer. The organic layer was washed with 5% hydrochloric acid and with brine, dried (MgSO_4) and evaporated to give a clear yellow liquid. Purification by flash chromatography and elution with 1:1 hexane/ethyl acetate containing 10% acetic acid yielded 2.0 g (28%) of benzyl[2,4-di(benzyloxycarbonyl)butyl]-phosphinic acid (3, $\text{R} = \text{CH}_2\text{Ph}$, $\text{R}_1 = \text{H}$) as a clear light yellow oil. R_f 0.37 (1:1 Hexane/EtOAc, 10% AcOH).

131

^1H NMR ($\text{d}_6\text{-DMSO}$): 7.2 ppm (m, 15H), 5.0 ppm (s, 4H), 3.0 (d, 2H), 2.8 ppm (m, 1H), 2.3 ppm (t, 2H), 1.9 ppm (m, 2H), 1.7 ppm (t, 1H).

5 2-[(Benzylhydroxyphosphinyl)methyl]pentanedioic acid

Benzyl[2,4-di(benzyloxycarbonyl)butyl]phosphinic acid (0.5 g, 1.0 mmol) in 20 mL of water containing 120 mg of 10% Pd/C was hydrogenated at 40 psi for 6 hours. Filtration through a Celite pad followed by evaporation
10 on high vacuum gave 0.17 g (57%) of 2-[(benzylhydroxyphosphinyl)methyl]-pentanedioic acid (4, R = CH_2Ph) as a white solid.

^1H NMR (D_2O): 7.1 ppm (m, 5H), 2.9 ppm (d, 2H), 2.4 ppm (m, 1H), 2.1 ppm (t, 2H), 1.8 ppm (m, 1H), 1.6 ppm (m,
15 3H).

Elemental Analysis

Calculated $\text{C}_{11}\text{H}_{17}\text{O}_6\text{P}$: C, 52.00; H, 5.71.

Found: C, 51.48; H, 5.70.

20

EXAMPLE 4

Preparation of 2-

[phenylethylhydroxyphosphinyl)methyl]pentanedioic acid

Scheme IV: R = $\text{CH}_2\text{CH}_2\text{Ph}$, $\text{R}_1 = \text{H}$

25 Phenethylphosphinic acid

Diethylchlorophosphite (15.6 g, 0.1 mol) in 100 mL

132

of dry diethyl ether was cooled to 5° C under an atmosphere of nitrogen. Phenethylmagnesium chloride (100 mL, 0.1 mol, 1.0 M in THF) was added dropwise over 2 hours while maintaining a temperature between 0-10° C.

5 A thick white slurry formed and stirred at room temperature overnight. The mixture was filtered under a nitrogen atmosphere and the filtrate evaporated under reduced pressure to give a clear and colorless liquid. The liquid was stirred as 15 mL of water was added

10 followed by 0.5 mL of concentrated hydrochloric acid. An exothermic reaction was observed and stirring continued for 15 minutes followed by extraction with ethyl acetate. The organics were combined, washed with brine, dried (MgSO₄) and evaporated. The clear liquid was brought up

15 in sodium hydroxide (40 mL, 2.0 M NaOH), stirred for 1 hour and washed once with diethyl ether. The aqueous layer was acidified to pH 1.0 with concentrated hydrochloric acid and extracted with ethyl acetate. The organics were combined, dried (MgSO₄) and evaporated to

20 give phenethylphosphinic acid (1, R = CH₂CH₂Ph, R₁ = H) (9.8 g, 58%) as a clear light yellow oil.

¹H NMR (d₆-DMSO): 7.2 ppm (m, 5H), 6.9 ppm (d, 1H), 2.8 ppm (m, 2H), 1.9 ppm (m, 2H).

25 2,4-Di(benzyloxycarbonyl)butyl(phenethyl)phosphinic acid
Phenethylphosphinic acid (1.0 g, 5.9 mmol) in 50 mL

133

of dry dichloromethane was cooled to -5°C under a nitrogen atmosphere. Triethylamine (2.3 g, 23 mmol) was added followed by trimethylsilyl chloride (2.2 g, 21 mmol) while the reaction temperature was maintained at 0°C . After 10 minutes dibenzyl 2-methylenepentane-dioate (2) (1.7 g, 5.2 mmol) in 10 mL of dichloromethane was added over 10 minutes. The reaction mixture was left to warm to room temperature and stirred overnight. The clear solution was cooled to 0°C and quenched with 5% hydrochloric acid followed by removal of the organic layer. The organic layer was washed with brine, dried (MgSO_4) and evaporated to give a clear light golden liquid. Purification by flash chromatography and elution with 1:1 Hexane/EtOAc containing 5% AcOH yielded 1.2 g (41%) of 2,4-di(benzyloxycarbonyl)-butyl(phenethyl)phosphinic acid (3, $\text{R} = \text{CH}_2\text{CH}_2\text{Ph}$, $\text{R}_1 = \text{H}$) as a clear and colorless oil.

^1H NMR (d_6 -DMSO): 7.2 ppm (m, 15H), 5.0 ppm (s, 4H), 3.3 ppm (m, 1H), 2.8 ppm (m, 4H), 2.3 ppm (m, 2H), 1.8 ppm (m, 4H).

2-[(Phenethylhydroxyphosphinyl)methyl]pentanedioic acid
2,4-Di(benzyloxycarbonyl)butyl(phenethyl)-phosphinic acid (1.1 g, 2.2 mmol) in 20 mL of water containing 120 mg of 10% Pd/C was hydrogenated at 40 psi overnight. Filtration through a Celite pad followed by evaporation

134

on high vacuum gave 0.8 g (114%) of 2-[(phenethylhydroxyphosphinyl)methyl]pentanedioic acid (4, R = CH₂CH₂Ph) as a white solid.

¹H NMR (D₂O): 7.2 ppm (m, 5H), 2.7 ppm (m, 2H), 2.5 ppm (m, 1H), 2.3 ppm (t, 2H), 1.9 ppm (m, 6H), 1.5 ppm (t, 1H)

Elemental Analysis

Calculated C₁₄H₁₉O₆P, 0.75H₂O, 0.5 AcOH: C, 50.35; H, 6.34.

Found: C, 50.26; H, 5.78.

10

EXAMPLE 5

Preparation of 2-[(3-

phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid

Scheme IV: R = CH₂CH₂CH₂Ph, R₁ = H

15

3-Phenylpropylphosphinic acid

To magnesium turnings (2.44 g, 0.10 mol) in 20 mL of dry diethyl ether under an atmosphere of nitrogen was added several iodine crystals. Phenylpropyl bromide (20.0 g, 0.10 mol) in 80 mL of diethyl ether was placed in a dropping funnel. Approximately 10 mL of the bromide solution was added to the magnesium turnings and stirring was initiated. After several minutes the iodine was consumed and additional phenylpropyl bromide was added while maintaining a temperature of 35° C. Once addition was complete (1.5 hours) the mixture was sealed and

stored at 5° C.

Diethylchlorophosphite (15.7 g, 0.1 mol) in 50 mL of dry diethyl ether was cooled to 5° C under an atmosphere of nitrogen. Phenethylmagnesium bromide (100 mL, 0.1 mol, 1.0 M solution of in Et₂O) was added dropwise over 2 hours while maintaining a temperature between 0 - 10° C. A thick white slurry formed and was stirred for an additional 30 minutes. The mixture was filtered under a nitrogen atmosphere and the filtrate evaporated under reduced pressure to give a clear and colorless liquid. To the liquid was added 20 mL of water followed by 0.5 ml of concentrated hydrochloric acid. An exothermic reaction was observed and stirring continued for 20 minutes followed by extraction with ethyl acetate. The organics were combined, washed with brine, dried (MgSO₄) and evaporated. To the clear liquid was added sodium hydroxide (40 mL, 2.0 M NaOH), the resulting solution stirred for 1 hour and then washed with diethyl ether. The aqueous layer was acidified to pH 1.0 with concentrated hydrochloric acid and extracted twice with ethyl acetate. The organics were combined, dried (MgSO₄) and evaporated to give 3-phenylpropylphosphinic acid (1, R = CH₂CH₂CH₂Ph, R₁ = H) (9.8 g, 53%) as a clear and colorless oil.

¹H NMR (d₆-DMSO): 7.2 ppm (m, 5H), 6.9 ppm (d, 1H), 2.6 ppm (t, 2H), 1.7 ppm (m, 2H), 1.6 ppm (m, 2H).

136

2,4-Di(benzyloxycarbonyl)butyl(3-phenylpropyl)-phosphinic acid

3-phenylpropylphosphinic acid (1.0 g, 5.4 mmol) in 50 mL of dry dichloromethane was cooled to -5° C under a nitrogen atmosphere. Triethylamine (2.2 g, 22 mmol) was added followed by trimethylsilyl chloride (2.1 g, 19 mmol) while the reaction temperature was maintained at 0° C. After 10 minutes dibenzyl 2-methylenepentanedioate (2) (1.6 g, 4.9 mmol) in 10 mL of dichloromethane was added over 10 minutes. The reaction mixture was warmed to room temperature and stirred overnight. The clear solution was cooled to 0° C and quenched with 5% hydrochloric acid followed by removal of the organic layer. The organic layer was washed with brine, dried (MgSO₄) and evaporated to give a clear yellow liquid. Purification by flash chromatography and elution with 4:1 hexane/ethyl acetate containing 5% acetic acid resulted in 1.5 g (56%) of 2,4-di(benzyloxycarbonyl)-butyl(3-phenylpropyl)phosphinic acid (3, R = CH₂CH₂CH₂Ph, R₁ = H) as a clear light yellow oil. *R_f* 0.58 (1:1 Hexane/EtOAc, 5% AcOH).

¹H NMR (d₆-DMSO): 7.2 ppm (m, 15H), 5.0 ppm (s, 4H), 2.7 ppm (m, 1H), 2.5 ppm (m, 5H), 2.2 ppm (m, 2H), 1.8 ppm (m, 3H), 1.6 ppm (m, 2H).

Elemental Analysis

Calculated C₂₉H₃₃O₆P, 1.3 H₂O: C, 65.48; H 6.75.

137

Found: C, 65.24; H, 6.39.

2-[(3-Phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid

5 2,4-Di(benzyloxycarbonyl)butyl(3-phenylpropyl)phosphinic acid (15) (1.4 g, 2.8 mmol) in 20 mL of water containing 150 mg of 10% Pd/C was hydrogenated at 40 psi overnight. Filtration through a Celite pad followed by evaporation on high vacuum gave
10 0.8 g (89%) of 2-[(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid (4, R = CH₂CH₂CH₂Ph) as a light yellow viscous oil.

¹H NMR (D₂O): 7.4 ppm (m, 5H), 2.7 ppm (m, 3H), 2.4 ppm (t, 3H), 1.8 ppm (m, 7H).

15 **Elemental Analysis**

Calculated C₁₅H₂₁O₆P, 0.75 H₂O, 0.75 AcOH: C, 51.23; H, 6.64.

Found: C, 50.85; H, 6.02.

20

EXAMPLE 6

Preparation of 2-[[4-methylbenzyl]hydroxyphosphinyl]methyl]pentanedioic acid

Scheme V: Compound 5, R = 4-methylbenzyl

Hexamethyldisilazane (21.1 mL, 100 mmol) was added to
25 vigorously stirred ammonium phosphinate (8.30 g, 100 mmol), and the resulting suspension was stirred at 105°

138

C for 2 hours. A solution of 4-methylbenzyl bromide (5.0 g, 27.0 mmol) was then added dropwise to the suspension at 0° C. The mixture was stirred at room temperature for 19 hours. The reaction mixture was then diluted with 5 dichloromethane (50 mL) and washed with 1 N HCl (50 mL). The organic layer was separated, dried over Na₂SO₄, and concentrated to give 4.72 g of a white solid. This was dissolved in dichloromethane (50 mL) and benzyl alcohol (3.24 g, 30 mmol) was added to the solution. 1,3-10 Dicyclohexylcarbodiimide (DCC)(6.19 g, 30 mmol) was then added to the solution at 0° C, and the suspension was stirred at room temperature for 14 hours. The solvent was removed under reduced pressure and the residue was suspended in EtOAc. The resulting suspension was 15 filtered and the filtrate was concentrated. The residue was purified by silica gel chromatography (hexanes: EtOAc, 4:1 to 1:1) to give 2.40 g of 4-methylbenzyl-O-benzylphosphinic acid (2, R = 4-methylbenzyl) as a white solid (34% yield). R_f 0.42 (EtOAc).

20 ¹H NMR (DMSO-d₆): δ 2.30 (s, 3H), 3.29 (d, 2H), 5.2 (m, 2H), 7.0 (d, 1H), 7.1-7.2 (m, 4H), 7.3-7.4 (m, 5H).

2,4-Di(benzyloxycarbonyl)-butyl(4-methylbenzyl)-o-benzylphosphinic acid

25 To a solution of 4-methylbenzyl-O-benzylphosphinic acid (2, R = 4-methylbenzyl) (2.16 g, 8.3 mmol) in THF (15

139

mL) was added sodium hydride (0.10 g, 60% dispersion in oil) followed by dibenzyl 2-methylenepentanedioate (3) (3.24 g) at 0° C, and the mixture was stirred at room temperature for 4 hours. The reaction mixture was then
5 diluted with EtOAc (50 mL) and poured into 1N HCl (50 mL). The organic layer was separated, dried over Na₂SO₄, and concentrated. This material was purified by silica gel chromatography (hexanes: EtOAc, 4:1 to 1:1) to give
3.41 g of 2,4-di(benzyloxycarbonyl)-butyl(4-
10 methylbenzyl)-o-benzylphosphinic acid (4, R = 4-methylbenzyl) as colorless oil (70% yield). R_f 0.61 (EtOAc).

¹H NMR (CDCl₃): δ 1.6-1.8 (m, 1H), 1.9-2.0 (m, 2H), 2.1-2.4 (m, 6H), 2.7-2.9 (m, 1H), 3.05 (dd, 2H), 4.8-5.1 (m,
15 6H), 7.0-7.1 (m, 4H), 7.2-7.4 (m, 15H).

2 - [[[4-Methylbenzyl)hydroxyphosphinyl]methyl] - pentanedioic acid

To a solution of 2,4-di(benzyloxycarbonyl)butyl(4-
20 methylbenzyl)-o-benzylphosphinic acid (0.70 g, 1.2 mmol) in ethanol (30 mL) was added Pd/C (5%, 0.10 g) and the suspension was shaken under hydrogen (50 psi) for 18 hours. The suspension was then filtered through a pad of Celite and concentrated under reduced pressure. The
25 resulting residue was dissolved in distilled water (5 mL), passed through a column of AG 50W-X8 resin (H⁺

. 140

form), and lyophilized to give 0.21 g of 2-[[[(4-methylbenzyl)hydroxyphosphinyl)methyl]-pentanedioic acid (5, R = 4-methylbenzyl) as a white solid (55% yield). R_f 0.62 (i-PrOH: H₂O, 7:3).

5 ¹H NMR (D₂O): δ 1.7-1.9 (m, 3H), 2.0-2.2 (m, 1H), 2.33 (dt, 7.4 Hz, 2H), 2.55-2.70 (m, 1H), 3.12 (d, 2H), 7.0-7.1 (m, 2H), 7.2-7.3 (m, 2H). Elemental Analysis
Calculated C₁₃H₁₇O₆P, 0.30 H₂O: C, 52.60; H, 6.18. Found: C, 52.60; H, 6.28.

10

EXAMPLE 7

Preparation of 2-[[[(4-Fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid

Scheme V: R = 4-fluorobenzyl

15 Prepared as described in the above example where R = methylbenzyl. R_f 0.64 (i-PrOH:H₂O, 7:3).

¹H NMR (D₂O): δ 1.7-1.9 (m, 3H), 2.0-2.2 (m, 1H), 2.3-2.4 (m, 2H), 2.55-2.70 (m, 1H), 3.12 (d, 2H), 7.0-7.1 (m, 2H), 7.2-7.3 (m, 2H).

20 Elemental Analysis

Calculated C₁₃H₁₆FO₆P, 0.25 H₂O: C, 48.38; H, 5.15. Found: C, 48.38; H, 5.15.

141

EXAMPLE 8

Preparation of 2-[[(4-Methoxybenzyl)hydroxyphosphinyl]
methyl]pentanedioic acid

Scheme V: R = 4-methoxybenzyl

5 Prepared as described in the above example where R = methylbenzyl. R_f 0.56 (*i*-PrOH: H₂O, 7:3).

¹H NMR (D₂O): δ 1.8-1.9 (m, 3H), 2.0-2.2 (m, 1H), 2.3-2.4 (m, 2H), 2.55-2.70 (m, 1H), 3.16 (d, 2H), 3.81 (s, 3H), 6.98 (d, 2H), 7.25 (d, 2H).

10 Elemental Analysis

Calculated C₁₄H₁₉O₇P, 0.30 H₂O: C, 50.09; H, 5.89.

Found: C, 49.98; H, 5.80.

EXAMPLE 9

15 Preparation of 2-[[(2-Fluorobenzyl)hydroxyphosphinyl]
methyl]pentanedioic acid

Scheme V: R = 2-fluorobenzyl)

Prepared as described in the above example where R = methylbenzyl. R_f 0.67 (*i*-PrOH: H₂O, 7:3).

20 ¹H NMR (D₂O): δ 1.8-1.9 (m, 3H), 2.0-2.2 (m, 1H), 2.3-2.4 (m, 2H), 2.55-2.70 (m, 1H), 3.28 (d, 2H), 7.1-7.5 (m, 4H).

Elemental Analysis

Calculated C₁₃H₁₆FO₆P, 0.10 H₂O: C, 48.79; H, 5.10. Found:

25 C, 48.84; H, 5.14.

142

EXAMPLE 10

Preparation of 2-[[[(Pentafluorobenzyl)hydroxyphosphinyl]
methyl]pentanedioic acid

5 Scheme V: R = pentafluorobenzyl

Prepared as described in the above example where R = methylbenzyl. R_f 0.69 (*i*-PrOH: H₂O, 7:3).

¹H NMR (D₂O): δ 1.8-2.0 (m, 3H), 2.1-2.3 (m, 1H), 2.3-2.5 (m, 2H), 2.7-2.9 (m, 1H), 3.29 (d, 2H).

10 Elemental Analysis

Calculated C₁₃H₁₂F₅O₆P, 0.45 H₂O: C, 39.20; H, 3.26. Found: C, 39.17; H, 3.28.

EXAMPLE 11

15 Preparation of 2-[(methylhydroxyphosphinyl)methyl]
pentanedioic acid

Scheme VI, Compound 9

2,4-Di(benzyloxycarbonyl)butylphosphinic acid (6)

20 Ammonium phosphinate (10 g, 0.12 mol) was placed in a round bottom flask with stirring under an atmosphere of nitrogen. Hexamethyldisilazane (HMDS, 25.5 mL, 0.12 mol) was added and the mixture heated to 110° C. After two hours the mixture was cooled to 0° C and dichloromethane
25 (120 ml) was added. After this was complete, dibenzyl-2-methylene pentanedioate (41 g, 0.13 mol) was added

143

dropwise. The mixture was allowed to warm to room temperature and stirred for 16 hours. The mixture was then quenched with 5% HCl (75 ml) and the organic layer removed. The organics were dried (MgSO₄) and evaporated under reduced pressure to give 42 g (90%) of a clear and colorless oil.

¹H NMR (CDCl₃): 7.36 ppm (m, 10H), 7.1 ppm (d, 1H), 5.19 ppm (s, 2H), 5.15 ppm (s, 2H), 2.92 ppm (m, 1H), 2.21 ppm (m, 6H).

10

2,4-Di(benzyloxycarbonyl)butylbenzylphosphinic acid (7)

To a solution of 2,4-di-(benzyloxycarbonyl)butylphosphinic acid (6) (19.3 g, 49.4 mmol) in tetrahydrofuran was added benzyl alcohol (5.3 g, 49.3 mmol) and dimethylamino pyridine (0.5 g). Dicyclohexylcarbodiimide (DCC, 12 g, 58 mmol) was added and a white precipitate formed. After 30 minutes the white suspension was filtered and the filtrate evaporated under reduced pressure. The clear and colorless oil was purified by flash chromatography and eluted with 1:1 Hexane/EtOAc to give 2,4-di(benzyloxycarbonyl)butylbenzylphosphinic acid (7) (11.5 g, 47%) as a clear and colorless oil. R_f 0.16 (1:1 Hexane/EtOAc).

¹H NMR (CDCl₃): 7.3 ppm (m, 15H), 7.2 ppm (d, 1H), 5.0 ppm (m, 6H), 2.9 ppm (m, 1H), 2.2 ppm (m, 3H), 1.9 ppm (m, 3H).

144

2,4-Di(benzyloxycarbonyl)butyl[hydroxy(phenyl)methyl]benzylphosphinic acid (8)

2,4-Di(benzyloxycarbonyl)butylbenzylphosphinic acid (7) in 5 mL of dry THF was added dropwise to a stirring cooled (0° C) mixture of sodium hydride (0.09 g, 2.3 mmol) in 15 mL of THF. After 15 minutes benzaldehyde (0.23 g, 2.2 mmol) was added via syringe while maintaining a temperature of 0° C. After 30 minutes the mixture was quenched with water and extracted with two portions of dichloromethane. The organics were combined and evaporated to give a clear colorless oil. The oil was chromatographed on silica and eluted with a 1:1 Hexane/EtOAc solvent system. The desired fractions were collected and evaporated to give 0.4 g (33%) of 2,4-di(benzyloxycarbonyl)butyl-[hydroxy(phenyl)methyl]benzylphosphinic acid (6) as a clear and colorless oil. R_f 0.18 (1:1 Hexane/EtOAc). 1H NMR ($CDCl_3$): 7.3 ppm (m, 20H), 5.2 ppm (m, 1H), 4.9 ppm (m, 6H), 2.8 ppm (dm, 1H), 2.2 ppm (m, 3H), 1.9 ppm (m, 3H).

2-([Hydroxy(phenyl)methyl]hydroxyphosphinylmethyl)-pentanedioic acid (9)

2,4-Di(benzyloxycarbonyl)butyl[hydroxy(phenyl)methyl]benzylphosphinic acid (6) (0.37 g, 0.6 mmol) in 25 mL of water containing 0.10 g of 10% Pd/C was

145

hydrogenated at 40 psi for 6 hours. The mixture was filtered through a pad of Celite and lyophilized to give 2-([hydroxy(phenyl)methyl]hydroxyphosphinyl-methyl)pentanedioic acid (9) (0.14 g, 70%) as a white solid.

^1H NMR (D_2O): 7.4 ppm (m, 5H), 5.0 ppm (d, 1H), 2.7 ppm (m, 1H), 2.4 ppm (m, 2H), 2.2 ppm (m, 1H), 1.9 ppm (m, 3H).

Elemental Analysis:

10 Calculated $\text{C}_{13}\text{H}_{17}\text{O}_7\text{P}$, 0.6 H_2O : C, 47.74; H, 5.61.

Found: C, 47.73; H, 5.68.

EXAMPLE 12

Preparation of dibenzyl 2-methylenepentanedioate using

15

Scheme III

Benzyl acrylate (500 g, 3.0 mol) was heated in an oil bath to 100° C. Heating was stopped and HMPT (10 g, 61 mmol) was added dropwise while maintaining an internal temperature below 140° C. Once addition was complete, 20 the mixture was stirred and cooled to room temperature. A slurry of silica (5:1 Hexane/EtOAc) was added and the mixture was placed in a column containing a plug of dry silica. The column was washed with 1:1 Hexane/EtOAc and the fractions were combined and evaporated to give 450 g 25 of clear light golden liquid. The liquid was distilled under high vacuum (200 μHg) at 185° C to give 212 g (42%)

146

of a clear and colorless liquid.

¹H NMR (CDCl₃): 7.3 ppm (m, 10H), 6.2 ppm (s, 1H), 5.6 ppm (s, 1H), 5.2 ppm (s, 2H), 5.1 ppm (s, 2H), 2.6 ppm (m, 4H).

5

EXAMPLE 13

Preparation of dibenzyl 2-

[[bis(benzyloxy)phosphoryl]methyl]pentanedioate using

Scheme III

10 Dibenzyl phosphite (9.5 g, 36 mmol) in 350 ml of dichloromethane was cooled to 0° C. To this stirring solution was added trimethyl aluminum (18.2 ml, 2.0 M solution in hexane, 36.4 mmol). After 30 minutes, dibenzyl 2-methylenepentanedioate (2) (6.0 g, 37 mmol) in
15 90 ml of dichloromethane was added dropwise over 10 minutes. The clear and colorless solution was then warmed to room temperature and left to stir overnight. The mixture was then quenched by the slow addition of 5% HCl. After stirring an additional 1.5 hours the lower
20 organic layer was removed and the aqueous layer extracted once with 100 ml of dichloromethane. The organics were combined, dried (MgSO₄), and evaporated to give a clear light golden liquid. The liquid was chromatographed on silica gel (4cm*30cm) and eluted with a gradient (4:1-
25 1:1) solvent system (Hexane/EtOAc). The fractions containing the desired product were combined and

147

evaporated to yield dibenzyl 2-[[bis(benzyloxy)-phosphoryl]methyl]pentanedioate (7.1 g, 42%) as a clear and colorless liquid. The liquid was then distilled on a Kughleror apparatus at 0.5 mm Hg and 195-200° C. The distillate was discarded and the remaining light golden oil was chromatographed on silica gel (1:1, Hexane/EtOAc) to give 2.9 g of dibenzyl 2-[[bis(benzyloxy)phosphoryl]methyl]pentanedioate as a clear and colorless oil. TLC R_f 0.5 (1:1 Hexane/EtOAc).

^1H NMR (CDCl_3): 7.1-7.4 (m, 20H), 5.05 (s, 2H), 4.8-5.03 (m, 6H), 2.8 (1H), 2.22-2.40 (m, 3H), 1.80-2.02 (m, 3H).

EXAMPLE 14

Preparation of 2-(phosphonomethyl)pentanedioic acid

(Compound 3) using Scheme III

Benzyl pentanedioate 2 (2.9 g, 4.9 mmol) was added to a mixture of 20 ml of methanol containing 0.29 g (6 mol %) of 10% Pd/C. This mixture was hydrogenated at 40 psi for 24 hours, filtered and evaporated to give 3 (1.0 g, 90%) as a clear slightly golden viscous oil.

^1H NMR (D_2O): 2.6-2.78 (m, 1H), 2.25-2.40 (m, 2H), 1.75-2.15 (m, 4H).

EXAMPLE 15

A patient is at risk of injury from an ischemic event. The patient may be pretreated with an effective

148

amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the pretreatment, the patient would be protected from any injury due to the ischemic event.

5

EXAMPLE 16

A patient is suffering from an ischemic event. The patient may be administered during or after the event, an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would recover or would not suffer any significant injury due to the ischemic event.

15

EXAMPLE 17

A patient has suffered injury from an ischemic event. The patient may be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would recover from the injury due to the ischemic event.

EXAMPLE 18

A patient is suffering from a glutamate abnormality. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition

149

of the present invention. It is expected that after the treatment, the patient would be protected from further injury due to the glutamate abnormality or would recover from the glutamate abnormality.

5

EXAMPLE 19

A patient is suffering from or has suffered from a nervous insult, such as that arising from a neurodegenerative disease or a neurodegenerative process.

10 The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from further injury due to the nervous insult or would recover from
15 the nervous insult.

EXAMPLE 20

A patient is suffering from Parkinson's disease. The patient may then be administered an effective amount
20 of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from further neurodegeneration or would recover from Parkinson's disease.

25

150

EXAMPLE 21

A patient is suffering from ALS. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from further neurodegeneration or would recover from ALS.

EXAMPLE 22

10 A patient is suffering from epilepsy. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from further neurodegeneration or would recover from epilepsy.

EXAMPLE 23

A patient is suffering from abnormalities in myelination/demyelination processes. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from further neurodegeneration or would recover from the abnormalities in myelination/demyelination processes.

151

EXAMPLE 24

A patient is suffering from or has suffered from a cerebrovascular accident, such as stroke. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from or would recover from any injury due to the cerebrovascular accident.

10

EXAMPLE 25

A patient is suffering from a head trauma. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from or would recover from any ischemic brain, spinal or peripheral injury resulting from the head trauma.

20

EXAMPLE 26

A patient is suffering from a spinal trauma. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from or would recover from any ischemic injury resulting from the

152

spinal trauma.

EXAMPLE 27

A patient is about to undergo surgery. The patient
5 may be administered an effective amount of a NAALADase
inhibitor or a pharmaceutical composition of the present
invention. It is expected that after the treatment, the
patient would not develop any ischemic brain, spinal or
peripheral injury resulting from or associated with the
10 surgery.

EXAMPLE 28

A patient is suffering from focal ischemia, such as
that associated with thromboembolytic occlusion of a
15 cerebral vessel, traumatic head injury, edema, or brain
tumors. The patient may then be administered an
effective amount of a NAALADase inhibitor or a
pharmaceutical composition of the present invention. It
is expected that after the treatment, the patient would
20 be protected from or would recover from any brain, spinal
or peripheral injury resulting from the focal ischemia.

EXAMPLE 29

A patient is suffering from global ischemia. The
25 patient may then be administered an effective amount of
a NAALADase inhibitor or a pharmaceutical composition of

153

the present invention. It is expected that after the treatment, the patient would be protected from or would recover from any brain, spinal or peripheral injury resulting from the global ischemia.

5

EXAMPLE 30

A patient is suffering from a cardiac arrest. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of
10 the present invention. It is expected that after the treatment, the patient would be protected from or would recover from any ischemic brain, spinal or peripheral injury associated with the cardiac arrest.

15

EXAMPLE 31

A patient is suffering from hypoxia, asphyxia or perinatal asphyxia. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It
20 is expected that after the treatment, the patient would be protected from or would recover from any ischemic brain, spinal or peripheral injury associated with the hypoxia, asphyxia or perinatal asphyxia.

25

EXAMPLE 32

A patient is suffering from a cerebro-cortical

154

injury. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would
5 be protected from or would recover from any ischemic brain injury resulting from the cerebro-cortical injury.

EXAMPLE 33

The patient is suffering from an injury to the
10 caudate nucleus. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from or would recover from any ischemic
15 brain injury resulting from the injury to the caudate nucleus.

EXAMPLE 34

A patient is diagnosed with a condition as
20 identified in these examples. An effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention may then be administered to the patient intravenously, intramuscularly, intraventricularly to the brain, rectally,
25 subcutaneously, intranasally, through a catheter with or without a pump, orally, through a transdermal patch,

155

topically, or through a polymer implant. After the treatment, the patient's condition would be expected to improve.

5

EXAMPLE 35

A patient is diagnosed with a condition as identified in these examples. A NAALADase inhibitor or a pharmaceutical composition of the present invention may then be administered to the patient in the form of a 100
10 mg/kg bolus, optionally followed by a 20 mg/kg per hour intravenous infusion over a two-hour period. After the treatment, the patient's condition would be expected to improve.

15

EXAMPLE 36

A patient is suffering from a cortical injury due to a condition identified in these examples. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of
20 the present invention. It is expected that after the treatment, the patient would be protected from further injury, or would exhibit at least 65% to at least 80% recovery from the cortical injury.

25

EXAMPLE 37

A patient is suffering from multiple sclerosis. The

156

patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from further demyelination or would recover from multiple sclerosis.

EXAMPLE 38


A patient is suffering from a peripheral neuropathy caused by Guillain-Barré syndrome. The patient may then be administered an effective amount of a NAALADase inhibitor or a pharmaceutical composition of the present invention. It is expected that after the treatment, the patient would be protected from further demyelination or would recover from the peripheral neuropathy.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modifications are intended to be included within the scope of the following claims.

WE CLAIM:

1. A pharmaceutical composition comprising:
 - (i) an effective amount of a NAALADase inhibitor
5 for treating a glutamate abnormality or
effecting a neuronal activity in an animal; and
 - (ii) a pharmaceutically acceptable carrier.
2. The pharmaceutical composition of claim 1,
10 further comprising at least one additional therapeutic agent.
3. The pharmaceutical composition of claim 1,
wherein said NAALADase inhibitor is selected from the
15 group consisting of a glutamate-derived hydroxyphosphinyl
derivative, an acidic peptide analog, a conformationally
restricted glutamate mimic and mixtures thereof.
4. The pharmaceutical composition of claim 3,
20 wherein the NAALADase inhibitor is an acidic peptide
analog selected from the group consisting of Asp-Glu,
Glu-Glu, Gly-Glu, gamma-Glu-Glu and Glu-Glu-Glu.
5. The pharmaceutical composition of claim 3,
25 wherein the NAALADase inhibitor is a glutamate-derived
hydroxyphosphinyl derivative of formula I:

158



5

or a pharmaceutically acceptable salt or hydrate thereof,
wherein:

R_1 is selected from the group consisting of hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl and Ar, wherein said R_1 is unsubstituted or substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_9 alkoxy, C_2 - C_9 alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

X is CR_3R_4 , O or NR_1 ;

R_3 and R_4 are independently selected from the group consisting of hydrogen, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, Ar, halo and mixtures thereof;

R_2 is selected from the group consisting of hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl and Ar, wherein said R_2 is unsubstituted or

substituted with carboxy, C₃-C₆ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

Ar is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuran-2-yl, tetrahydropyran-2-yl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, benzyl and phenyl, wherein said Ar is unsubstituted or substituted with halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino or mixtures thereof.

15

6. The pharmaceutical composition of claim 5, wherein X is CH₂.

7. The pharmaceutical composition of claim 6, wherein R₂ is substituted with carboxy.

8. The pharmaceutical composition of claim 7, wherein:

R₁ is hydrogen, C₁-C₄ straight or branched chain alkyl, C₂-C₄ straight or branched chain alkenyl, C₃-C₆ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl or phenyl, wherein

160

said R_1 is unsubstituted or substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 -
5 C_4 alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof; and

R_2 is C_1 - C_2 alkyl.

9. The pharmaceutical composition of claim 8,
10 wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 2-(phosphonomethyl)pentanedioic acid;
 - 2-(phosphonomethyl)succinic acid;
 - 2-[[(2-carboxyethyl)hydroxyphosphinyl]methyl]pentanedioic
15 acid;
 - 2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 20 2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[(cyclohexyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
 - 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[benzylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 25 2-[[(phenylmethyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

161

- 2- [[(phenylethyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2- [[(phenylpropyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 5 2- [[(phenylbutyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2- [[(4-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2- [[(4-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic
- 10 acid;
- 2- [[(2-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2- [[(pentafluorobenzyl) hydroxyphosphinyl] methyl] pentane-dioic acid;
- 15 2- [[(methoxybenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2- [[(2,3,4-trimethoxyphenyl) hydroxyphosphinyl] methyl] -pentanedioic acid;
- 2- [[(phenylprop-2-enyl) hydroxyphosphinyl] methyl] pentane-
- 20 dioic acid;
- 2- [[(2-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2- [[((hydroxy) phenylmethyl) hydroxyphosphinyl] methyl] -pentanedioic acid;
- 25 2- [[(3-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;

- 2-[[(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]-pentanedioic acid; and
5 pharmaceutically acceptable salts and hydrates thereof.

10. The pharmaceutical composition of claim 9, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 10 2-(phosphonomethyl)pentanedioic acid;
2-(phosphonomethyl)succinic acid;
2-[[(2-carboxyethyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
15 2-[(phenylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[((hydroxy)phenylmethyl)hydroxyphosphinyl)methyl]-pentanedioic acid;
2-[(butylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[(3-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic
20 acid;
2-[(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic acid;
25 2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[(phenylethylhydroxyphosphinyl)methyl]pentanedioic

163

- acid;
2- [[(4-methylbenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2- [[(4-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
5 acid;
2- [[(4-methoxybenzyl)hydroxyphosphinyl]methyl]pentane-
dioic acid;
2- [[(3-trifluoromethylbenzyl)hydroxyphosphinyl]methyl] -
pentanedioic acid;
10 2- [[(2-fluorobenzyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2- [[(pentafluorobenzyl)hydroxyphosphinyl]methyl]pentane-
dioic acid;
2- [[(phenylprop-2-enyl)hydroxyphosphinyl]methyl]pentane-
15 dioic acid;
2- [[(aminomethyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
2- [[(aminoethyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
20 2- [[(aminopropyl)hydroxyphosphinyl]methyl]pentanedioic
acid; and
pharmaceutically acceptable salts and hydrates thereof.

11. The pharmaceutical composition of claim 10,
25 wherein the glutamate-derived hydroxyphosphinyl
derivative is 2-(phosphonomethyl)pentanedioic acid or a

pharmaceutically acceptable salt or hydrate thereof.

12. The pharmaceutical composition of claim 7, wherein:

5 R_1 is hydrogen, C_1 - C_4 straight or branched chain alkyl, C_2 - C_4 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, benzyl or phenyl, wherein said R_1 is unsubstituted or substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro,
10 trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 - C_4 alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof; and

R_2 is C_3 - C_9 alkyl.

15

13. The pharmaceutical composition of claim 12, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
- 20 2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
- 25 2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;

2-[(methylhydroxyphosphinyl)methyl]decanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid; and
pharmaceutically acceptable salts and hydrates thereof.

5 14. The pharmaceutical composition of claim 7,
wherein R₁ is 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl,
3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl,
3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl or C₁-C₄
straight or branched chain alkyl substituted with 2-
10 indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl,
tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-
pyridyl or 4-pyridyl.

15 15. The pharmaceutical composition of claim 14,
wherein the glutamate-derived hydroxyphosphinyl
derivative is selected from the group consisting of:
2-[[[(2-pyridyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
2-[[[(3-pyridyl)methylhydroxyphosphinyl)methyl]pentane-
20 dioic acid;
2-[[[(4-pyridyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
2-[[[(3-pyridyl)ethylhydroxyphosphinyl)methyl]pentane-
dioic acid;
25 2-[[[(3-pyridyl)propylhydroxyphosphinyl)methyl]pentane-
dioic acid;

- 2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl)methyl]-
pentanedioic acid;
- 2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl)methyl]-
pentanedioic acid;
- 5 2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl)methyl]-
pentanedioic acid;
- 2-[[(2-tetrahydropyranyl)hydroxyphosphinyl)methyl]-
pentanedioic acid;
- 2-[[(3-tetrahydropyranyl)hydroxyphosphinyl)methyl]-
10 pentanedioic acid;
- 2-[[(4-tetrahydropyranyl)hydroxyphosphinyl)methyl]-
pentanedioic acid;
- 2-[[(2-indolyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 15 2-[[(3-indolyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[(4-indolyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[(3-indolyl)ethylhydroxyphosphinyl)methyl]pentane-
20 dioic acid;
- 2-[[(3-indolyl)propylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[(2-thienyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 25 2-[[(3-thienyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;

167

- 2-[[[(4-thienyl)methylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 5 2-[[[(3-thienyl)propylhydroxyphosphinyl)methyl]pentane-
dioic acid;
- 2-[[[(2-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 2-[[[(3-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
10 acid;
- 2-[[[(4-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl)methyl]pentane-
dioic acid;
- 15 2-[[[(2-indolyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 2-[[[(3-indolyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 2-[[[(4-indolyl)hydroxyphosphinyl)methyl]pentanedioic
20 acid;
- 2-[[[(2-thienyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 2-[[[(3-thienyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 25 2-[[[(4-thienyl)hydroxyphosphinyl)methyl]pentanedioic
acid; and

pharmaceutically acceptable salts and hydrates thereof.

16. The pharmaceutical composition of claim 7, wherein R₁ is 1-naphthyl, 2-naphthyl, or C₁-C₄ straight or
5 branched chain alkyl substituted with 1-naphthyl or 2-naphthyl.

17. The pharmaceutical composition of claim 16, wherein the glutamate-derived hydroxyphosphinyl
10 derivative is selected from the group consisting of:
2-[[[(1-naphthyl)hydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl)methyl]pentanedioic acid;
15 2-[[[(1-naphthyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl)methyl]pentane-
20 dioic acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl)methyl]pentanedioic acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl)methyl]pentanedioic acid;
25 2-[[[(2-naphthyl)propylhydroxyphosphinyl)methyl]pentanedioic acid;

2-[[[(1-naphthyl)butylhydroxyphosphinyl)methyl]pentane-dioic acid;

2-[[[(2-naphthyl)butylhydroxyphosphinyl)methyl]pentane-dioic acid; and

5 pharmaceutically acceptable salts and hydrates thereof.

18. The pharmaceutical composition of claim 6, wherein:

R₂ is selected from the group consisting of
10 hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl and phenyl, wherein said R₂ is unsubstituted or substituted with C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆
15 straight or branched chain alkenyl, C₁-C₄ alkoxy, phenyl or mixtures thereof.

19. The pharmaceutical composition of claim 18, wherein:

20 R₁ is hydrogen, C₁-C₄ straight or branched chain alkyl, C₂-C₄ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl or phenyl, wherein said R₁ is unsubstituted or substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro,
25 trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-

C, alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof.

20. The pharmaceutical composition of claim 19,
5 wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 3-(methylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(propylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 10 3-(butylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(cyclohexylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-((cyclohexyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(phenylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 15 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(phenylethylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(phenylpropylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(phenylbutylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-((2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl)-2-phenylpropanoic acid;
 - 20 3-(phenylprop-2-enylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-(benzylhydroxyphosphinyl)-2-ethylpropanoic acid;
 - 3-(benzylhydroxyphosphinyl)-2-propylpropanoic acid;
 - 25 3-(benzylhydroxyphosphinyl)-2-butylpropanoic acid;
 - 3-(benzylhydroxyphosphinyl)-2-cyclohexylpropanoic acid;

171

- 3- (benzylhydroxyphosphinyl) -2- (cyclohexyl) methylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- phenylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- benzylpropanoic acid;
- 5 3- (benzylhydroxyphosphinyl) -2- phenylethylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- phenylpropylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- phenylbutylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- (2,3,4-trimethoxyphenyl) -propanoic acid;
- 10 3- (benzylhydroxyphosphinyl) -2- phenylprop-2-enylpropanoic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

21. The pharmaceutical composition of claim 6,

15 wherein at least one of R₁ and R₂ is 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or C₁-C₄ straight or branched chain alkyl substituted with 2-indolyl 3-indolyl, 4-indolyl, 2-

20 furyl, 3-furyl, tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl or 4-pyridyl.

22. The pharmaceutical composition of claim 21,

wherein the glutamate-derived hydroxyphosphinyl

25 derivative is selected from the group consisting of:

3- [(2-pyridyl) methylhydroxyphosphinyl] -2- phenylpropanoic

172

- acid;
- 3-[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 5 acid;
- 3-[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 10 3-[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 15 acid;
- 3-[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 20 3-[(4-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 25 acid;
- 3-[(2-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic

- acid;
3-[(3-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(4-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic
5 acid;
3-[(3-thienyl)ethylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-thienyl)propylhydroxyphosphinyl]-2-phenylpropanoic
acid;
10 3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)methylpropanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)methylpropanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)methylpropanoic
15 acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)ethylpropanoic
acid;
3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propylpropanoic
acid;
20 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)methyl-
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)ethyl-
propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propyl-
25 propanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2-indolyl)methylpropanoic

174

- acid;
- 3- (benzylhydroxyphosphinyl) -2- (3-indolyl) methylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- (4-indolyl) methylpropanoic
- 5 acid;
- 3- (benzylhydroxyphosphinyl) -2- (3-indolyl) ethylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- (3-indolyl) propylpropanoic acid;
- 10 3- (benzylhydroxyphosphinyl) -2- (2-thienyl) methylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- (3-thienyl) methylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- (4-thienyl) methylpropanoic
- 15 acid;
- 3- (benzylhydroxyphosphinyl) -2- (3-thienyl) ethylpropanoic acid;
- 3- (benzylhydroxyphosphinyl) -2- (3-thienyl) propylpropanoic acid; and
- 20 pharmaceutically acceptable salts and hydrates thereof.

23. The pharmaceutical composition of claim 18, wherein R₁ is 1-naphthyl, 2-naphthyl, or C₁-C₄ straight or branched chain alkyl substituted with 1-naphthyl or 2-

25 naphthyl.

24. The pharmaceutical composition of claim 23, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 3-((1-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
 - 5 3-((2-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-((1-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-((2-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 10 3-((1-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-((2-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-((1-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 15 3-((2-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 3-((1-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid;
 - 20 3-((2-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

25. The pharmaceutical composition of claim 5, wherein X is O.

26. The pharmaceutical composition of claim 25, wherein R₂ is substituted with carboxy.

27. The pharmaceutical composition of claim 26,
5 wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 10 2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 15 2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
 - 20 2-[[4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
 - 2-[[2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
 - 25 2-[[pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;

177

- 2- [[(methoxybenzyl) hydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(2,3,4-trimethoxyphenyl) hydroxyphosphinyl] oxy] - pentanedioic acid;
- 5 2- [[(1-naphthyl) hydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(2-naphthyl) hydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(1-naphthyl) methylhydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(2-naphthyl) methylhydroxyphosphinyl] oxy] pentanedioic
- 10 acid;
- 2- [[(1-naphthyl) ethylhydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(2-naphthyl) ethylhydroxyphosphinyl] oxy] pentanedioic acid;
- 15 2- [[(1-naphthyl) propylhydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(2-naphthyl) propylhydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(1-naphthyl) butylhydroxyphosphinyl] oxy] pentanedioic
- 20 acid;
- 2- [[(2-naphthyl) butylhydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[(phenylprop-2-enyl) hydroxyphosphinyl] oxy] pentanedioic acid;
- 25 2- [[(benzylhydroxyphosphinyl] oxy] pentanedioic acid;
- 2- [[((hydroxy) phenylmethyl) hydroxyphosphinyl] oxy] pentane-

- dioic acid;
- 2-[[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 5 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-(phosphono)oxy]pentanedioic acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy]-
- 10 pentanedioic acid;
- 2-[[methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 15 2-[[cyclohexylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 25 2-[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;

- 2-[[[(2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 5 2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 10 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 15 2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 20 2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 25 2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;

- 2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;
5 2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;
2-[(methylhydroxyphosphinyl)oxy]decanedioic acid;
2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-methylethanoic acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-ethylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-propylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-butylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-cyclohexylethanoic
acid;
15 2-[[benzylhydroxyphosphinyl]oxy]-2-(cyclohexyl)methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-benzylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethylethanoic
20 acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylpropylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylbutylethanoic
acid;
25 2-[[benzylhydroxyphosphinyl]oxy]-2-(2,3,4-trimethoxy-
phenyl)ethanoic acid;

181

- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethyl-
- 10 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propyl-ethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butyl-
- 20 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enyl-ethanoic acid;
- 2-[[(2-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 25 2-[[(3-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

SUBSTITUTE SHEET (RULE 26)

182

- 2-[[(4-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 5 2-[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy] - pentanedioic acid;
- 2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy] -
- 10 pentanedioic acid;
- 2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy] - pentanedioic acid;
- 2-[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 15 2-[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic
- 20 acid;
- 2-[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 25 2-[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;

SUBSTITUTE SHEET (RULE 26)

- 2-[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
2-[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
5 2-[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic acid; and
pharmaceutically acceptable salts and hydrates thereof.

28. The pharmaceutical composition of claim 25,
10 wherein:

R₂ is selected from the group consisting of hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl and phenyl, wherein said R₂ is
15 unsubstituted or substituted with C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, phenyl or mixtures thereof.

- 20 29. The pharmaceutical composition of claim 28, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
2-[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
25 2-[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;

- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 5 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 10 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 15 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 20 2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 25 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;

- 2-[[(4-thienyl) methylhydroxyphosphinyl] oxy] -2-phenyl-ethanoic acid;
- 2-[[(3-thienyl) ethylhydroxyphosphinyl] oxy] -2-phenyl-ethanoic acid;
- 5 2-[[(3-thienyl) propylhydroxyphosphinyl] oxy] -2-phenyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(2-pyridyl) methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(3-pyridyl) methyl-
- 10 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(4-pyridyl) methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(3-pyridyl) ethyl-ethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl] oxy] -2-(3-pyridyl) propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(tetrahydrofuranyl) -methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(tetrahydrofuranyl) -
- 20 ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(tetrahydrofuranyl) -propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl] oxy] -2-(2-indolyl) methyl-ethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl] oxy] -2-(3-indolyl) methyl-ethanoic acid;

- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (4-indolyl)methyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-indolyl)ethyl-ethanoic acid;
- 5 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-indolyl)propyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (2-thienyl)methyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-thienyl)methyl-ethanoic acid;
- 10 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (4-thienyl)methyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-thienyl)ethyl-ethanoic acid;
- 15 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-thienyl)propyl-ethanoic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

30. The pharmaceutical composition of claim 5,
20 wherein X is NR_1 .

31. The pharmaceutical composition of claim 30,
wherein R_2 is substituted with carboxy.

25 32. The pharmaceutical composition of claim 31,
wherein the glutamate-derived hydroxyphosphinyl

derivative is selected from the group consisting of:

- 2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[cyclohexyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;
- 10 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;
- 15 2-[[4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 20 2-[[pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 25 2-[[2,3,4-trimethoxyphenyl)hydroxyphosphinyl]amino]pentanedioic acid;

- 2-[[[(1-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 5 2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 10 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 15 2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-naphthyl)butylhydroxyphosphinyl]amino]pentanedioic acid;
- 20 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]-2-pentanedioic acid;
- 25 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]amino]-

- pentanedioic acid;
- 2-[[[(3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic acid;
- 5 acid;
- 2-[(phosphono)amino]pentanedioic acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]amino]-pentanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]hexanedioic acid;
- 10 2-[(benzylhydroxyphosphinyl)amino]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]octanedioic acid;
- 15 2-[(methylhydroxyphosphinyl)amino]nonanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]nonanedioic acid;
- 2-[(methylhydroxyphosphinyl)amino]decanedioic acid;
- 2-[(benzylhydroxyphosphinyl)amino]decanedioic acid;
- 3-[[[(2-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 20 acid;
- 3-[[[(3-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[[(4-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 25 3-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

- 3- [[(3-pyridyl) propylhydroxyphosphinyl] amino] pentanedioic acid;
- 3- [[(tetrahydrofuranyl) methylhydroxyphosphinyl] amino] - pentanedioic acid;
- 5 3- [[(tetrahydrofuranyl) ethylhydroxyphosphinyl] amino] - pentanedioic acid;
- 3- [[(tetrahydrofuranyl) propylhydroxyphosphinyl] amino] - pentanedioic acid;
- 3- [[(2-indolyl) methylhydroxyphosphinyl] amino] pentanedioic
- 10 acid;
- 3- [[(3-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 3- [[(4-indolyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 15 3- [[(3-indolyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 3- [[(3-indolyl) propylhydroxyphosphinyl] amino] pentanedioic acid;
- 3- [[(2-thienyl) methylhydroxyphosphinyl] amino] pentanedioic
- 20 acid;
- 3- [[(3-thienyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 3- [[(4-thienyl) methylhydroxyphosphinyl] amino] pentanedioic acid;
- 25 3- [[(3-thienyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;

3-[[[(3-thienyl)propylhydroxyphosphinyl]amino]pentanedioic acid; and
pharmaceutically acceptable salts and hydrates thereof.

5 33. The pharmaceutical composition of claim 30, wherein R₂ is selected from the group consisting of hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl and phenyl, wherein said R₂ is
10 unsubstituted or substituted with C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl, C₁-C₈ alkoxy, phenyl or mixtures thereof.

15 34. The pharmaceutical composition of claim 33, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
2-[[methylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[ethylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
20 2-[[propylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[butylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[cyclohexylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]-2-phenyl-
25 ethanoic acid;
2-[[phenylhydroxyphosphinyl]amino]-2-phenylethanoic acid;

- 2-[[benzylhydroxyphosphinyl]amino]-2-phenylethanoic acid;
2-[[phenylethylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
2-[[phenylpropylhydroxyphosphinyl]amino]-2-phenylethanoic
5 acid;
2-[[phenylbutylhydroxyphosphinyl]amino]-2-phenylethanoic
acid;
2-[[[(2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl]-amino]-
2-phenylethanoic acid;
10 2-[[[(1-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic
acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]amino]-2-phenylethanoic
acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]-2-phenyl-
15 ethanoic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
20 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(2-naphthyl)propylhydroxyphosphinyl]amino]-2-phenyl-
25 ethanoic acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]amino]-2-phenyl-

- ethanoic acid;
2-[[(2-naphthyl) butylhydroxyphosphinyl] amino] -2-phenyl-
ethanoic acid;
2-[[phenylprop-2-enylhydroxyphosphinyl] amino] -2-phenyl-
5 ethanoic acid;
2-[[benzylhydroxyphosphinyl] amino] -2-methylethanoic acid;
2-[[benzylhydroxyphosphinyl] amino] -2-ethylethanoic acid;
2-[[benzylhydroxyphosphinyl] amino] -2-propylethanoic acid;
2-[[benzylhydroxyphosphinyl] amino] -2-butylethanoic acid;
10 2-[[benzylhydroxyphosphinyl] amino] -2-cyclohexylethanoic
acid;
2-[[benzylhydroxyphosphinyl] amino] -2-(cyclohexyl) methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl] amino] -2-phenylethanoic acid;
15 2-[[benzylhydroxyphosphinyl] amino] -2-benzylethanoic acid;
2-[[benzylhydroxyphosphinyl] amino] -2-phenylethylethanoic
acid;
2-[[benzylhydroxyphosphinyl] amino] -2-phenylpropylethanoic
acid;
20 2-[[benzylhydroxyphosphinyl] amino] -2-phenylbutylethanoic
acid;
2-[[benzylhydroxyphosphinyl] amino] -2-(2,3,4-trimethoxy-
phenyl)ethanoic acid;
2-[[benzylhydroxyphosphinyl] amino] -2-(1-naphthyl) ethanoic
25 acid;
2-[[benzylhydroxyphosphinyl] amino] -2-(2-naphthyl) ethanoic

- acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)methyl-
- 5 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)ethyl-ethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(1-naphthyl)butyl-
- 15 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-(2-naphthyl)butyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]amino]-2-phenolprop-2-enyl-ethanoic acid;
- 20 2-[[[(2-pyridyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]amino]-2-phenyl-ethanoic acid;
- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]amino]-2-phenyl-
- 25 ethanoic acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]amino]-2-phenyl-

- ethanoic acid;
2-[[[(3-pyridyl)propylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino]-2-
5 phenylethanoic acid;
2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino]-2-
phenylethanoic acid;
2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino]-2-
phenylethanoic acid;
10 2-[[[(2-indolyl)methylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(3-indolyl)methylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(4-indolyl)methylhydroxyphosphinyl]amino]-2-phenyl-
15 ethanoic acid;
2-[[[(3-indolyl)ethylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(3-indolyl)propylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
20 2-[[[(2-thienyl)methylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(3-thienyl)methylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[[(4-thienyl)methylhydroxyphosphinyl]amino]-2-phenyl-
25 ethanoic acid;
2-[[[(3-thienyl)ethylhydroxyphosphinyl]amino]-2-phenyl-

- ethanoic acid;
2-[[(3-thienyl)propylhydroxyphosphinyl]amino]-2-phenyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-pyridyl)methyl-
5 ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-pyridyl)methyl-
ethanoic acid;
10 2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)ethyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-pyridyl)propyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)-
15 methylethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)-
ethylethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(tetrahydrofuranyl)-
propylethanoic acid;
20 2-[[benzylhydroxyphosphinyl]amino]-2-(2-indolyl)methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-indolyl)methyl-
25 ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)ethyl-

ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-indolyl)propyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(2-thienyl)methyl-
5 ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)methyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(4-thienyl)methyl-
ethanoic acid;
10 2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)ethyl-
ethanoic acid;
2-[[benzylhydroxyphosphinyl]amino]-2-(3-thienyl)propyl-
ethanoic acid; and
pharmaceutically acceptable salts and hydrates thereof.

15

35. A method of treating a glutamate abnormality in an animal, comprising administering an effective amount of a NAALADase inhibitor to said animal.

20 36. The method of claim 35, wherein the glutamate abnormality is selected from the group consisting of epilepsy, stroke, Alzheimer's disease, Parkinson's Disease, Amyotrophic Lateral Sclerosis (ALS), Huntington's Disease, schizophrenia, chronic pain,
25 ischemia, peripheral neuropathy, traumatic brain injury and physical damage to the spinal cord.

198

37. The method of claim 36, wherein the glutamate abnormality is stroke.

38. The method of claim 36, wherein the glutamate
5 abnormality is Parkinson's Disease.

39. The method of claim 36, wherein the glutamate abnormality is Amyotrophic Lateral Sclerosis (ALS).

10 40. The method of claim 36, wherein the glutamate abnormality is ischemic spinal cord injury.

41. The method of claim 35, wherein the NAALADase inhibitor is administered in combination with at least
15 one additional therapeutic agent.


42. The method of claim 35, wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative, an acidic peptide analog or a mixture
20 thereof.

43. The method of claim 42, wherein the NAALADase inhibitor is an acidic peptide analog selected from the group consisting of Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-
25 Glu and Glu-Glu-Glu.

SUBSTITUTE SHEET (RULE 26)

199

44. The method of claim 42, wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative of formula I:



or a pharmaceutically acceptable salt or hydrate thereof,
10 wherein:

R_1 is selected from the group consisting of hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl and Ar, wherein said R_1 is unsubstituted or
15 substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_9 alkoxy, C_2 - C_9 alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

20 X is CR_3R_4 , O or NR_1 ;

R_3 and R_4 are independently selected from the group consisting of hydrogen, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, Ar, halo and mixtures
25 thereof;

R_2 is selected from the group consisting of

SUBSTITUTE SHEET (RULE 26)

200

hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl and Ar, wherein said R₂ is unsubstituted or substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino, Ar or mixtures thereof;

Ar is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuran-2-yl, tetrahydropyran-2-yl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, benzyl and phenyl, wherein said Ar is unsubstituted or substituted with halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino or mixtures thereof.

45. The method of claim 44, wherein X is CH₂.

20

46. The method of claim 45, wherein R₂ is substituted with carboxy.

47. The method of claim 46, wherein:

R₁ is hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, C₃-C₈

201

cycloalkyl, C₅-C₇ cycloalkenyl, benzyl or phenyl, wherein said R₁ is unsubstituted or substituted with carboxy, C₁-C₆ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof; and

R₂ is C₁-C₂ alkyl.

- 10 48. The method of claim 47, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 2-(phosphonomethyl)pentanedioic acid;
 - 2-(phosphonomethyl)succinic acid;
 - 15 2-[[(2-carboxyethyl)hydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[methylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[ethylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[propylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 20 2-[[butylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[cyclohexylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[(cyclohexyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 2-[[phenylhydroxyphosphinyl]methyl]pentanedioic acid;
 - 25 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
 - 2-[[(phenylmethyl)hydroxyphosphinyl]methyl]pentanedioic

SUBSTITUTE SHEET (RULE 26)

- acid;
2-[[(phenylethyl) hydroxyphosphinyl] methyl] pentanedioic
acid;
2-[[(phenylpropyl) hydroxyphosphinyl] methyl] pentanedioic
5 acid;
2-[[(phenylbutyl) hydroxyphosphinyl] methyl] pentanedioic
acid;
2-[[(4-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic
acid;
10 2-[[(4-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic
acid;
2-[[(2-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic
acid;
2-[[(pentafluorobenzyl) hydroxyphosphinyl] methyl] pentane-
15 dioic acid;
2-[[(methoxybenzyl) hydroxyphosphinyl] methyl] pentanedioic
acid;
2-[[(2,3,4-trimethoxyphenyl) hydroxyphosphinyl] methyl] -
pentanedioic acid;
20 2-[[(phenylprop-2-enyl) hydroxyphosphinyl] methyl] pentane-
dioic acid;
2-[[(2-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic
acid;
2-[[((hydroxy) phenylmethyl) hydroxyphosphinyl] methyl] -
25 pentanedioic acid;
2-[[(3-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic

SUBSTITUTE SHEET (RULE 26)

acid;

2-[[[(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]-
5 pentanedioic acid; and
pharmaceutically acceptable salts and hydrates thereof.

49. The method of claim 48, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the
10 group consisting of:

2-(phosphonomethyl)pentanedioic acid;

2-(phosphonomethyl)succinic acid;

2-[[[(2-carboxyethyl)hydroxyphosphinyl)methyl]pentanedioic acid;

15 2-[(benzylhydroxyphosphinyl)methyl]pentanedioic acid;

2-[(phenylhydroxyphosphinyl)methyl]pentanedioic acid;

2-[[[(hydroxy)phenylmethyl)hydroxyphosphinyl)methyl]-pentanedioic acid;

2-[(butylhydroxyphosphinyl)methyl]pentanedioic acid;

20 2-[[[(3-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic acid;

2-[[[(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic
25 acid;

2-[(methylhydroxyphosphinyl)methyl]pentanedioic acid;

204

2-[(phenylethylhydroxyphosphinyl)methyl]pentanedioic acid;

2-[(4-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

5 2-[(4-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[(4-methoxybenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

10 2-[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[(pentafluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;

15 2-[(phenylprop-2-enyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[(aminomethyl)hydroxyphosphinyl)methyl]pentanedioic acid;

20 2-[(aminoethyl)hydroxyphosphinyl)methyl]pentanedioic acid;

2-[(aminopropyl)hydroxyphosphinyl)methyl]pentanedioic acid; and

pharmaceutically acceptable salts and hydrates thereof.

25 50. The method of claim 49, wherein the glutamate-derived hydroxyphosphinyl derivative is 2-

205

(phosphonomethyl)pentanedioic acid or
a pharmaceutically acceptable salt or hydrate thereof.

51. The method of claim 46, wherein:

5 R_1 is hydrogen, C_1 - C_4 straight or branched chain
alkyl, C_2 - C_6 straight or branched chain alkenyl, C_3 - C_8
cycloalkyl, C_5 - C_7 cycloalkenyl, benzyl or phenyl, wherein
said R_1 is unsubstituted or substituted with carboxy, C_3 -
 C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro,
10 trifluoromethyl, C_1 - C_6 straight or branched chain alkyl,
 C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 -
 C_4 alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl
or mixtures thereof; and

R_2 is C_3 - C_9 alkyl.

15

52. The method of claim 51, wherein the glutamate-
derived hydroxyphosphinyl derivative is selected from the
group consisting of:

2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
20 2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
25 2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;

SUBSTITUTE SHEET (RULE 26)

206

2-[(methylhydroxyphosphinyl)methyl]decanedioic acid;
2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid; and
pharmaceutically acceptable salts and hydrates thereof

5 53. The method of claim 46, wherein R₁ is 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl or C₁-C₄ straight or branched chain alkyl substituted with 2-indolyl, 3-
10 indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl or 4-pyridyl.

54. The method of claim 53, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the
15 group consisting of:

2-[[[(2-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
20 2-[[[(4-pyridyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-pyridyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
2-[[[(3-pyridyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
25 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]methyl]-

SUBSTITUTE SHEET (RULE 26)

- pentanedioic acid;
2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]methyl]-
pentanedioic acid;
2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]methyl]-
5 pentanedioic acid;
2-[[[(2-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
pentanedioic acid;
2-[[[(3-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
pentanedioic acid;
10 2-[[[(4-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
pentanedioic acid;
2-[[[(2-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
2-[[[(3-indolyl)methylhydroxyphosphinyl]methyl]pentane-
15 dioic acid;
2-[[[(4-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
2-[[[(3-indolyl)ethylhydroxyphosphinyl]methyl]pentane-
dioic acid;
20 2-[[[(3-indolyl)propylhydroxyphosphinyl]methyl]pentane-
dioic acid;
2-[[[(2-thienyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
2-[[[(3-thienyl)methylhydroxyphosphinyl]methyl]pentane-
25 dioic acid;
2-[[[(4-thienyl)methylhydroxyphosphinyl]methyl]pentane-

- dioic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl)methyl]pentane-
- dioic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl)methyl]pentane-
- 5 dioic acid;
- 2-[[[(2-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
- acid;
- 2-[[[(3-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
- acid;
- 10 2-[[[(4-pyridyl)hydroxyphosphinyl)methyl]pentanedioic
- acid;
- 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl)methyl]pentane-
- dioic acid;
- 2-[[[(2-indolyl)hydroxyphosphinyl)methyl]pentanedioic
- 15 acid;
- 2-[[[(3-indolyl)hydroxyphosphinyl)methyl]pentanedioic
- acid;
- 2-[[[(4-indolyl)hydroxyphosphinyl)methyl]pentanedioic
- acid;
- 20 2-[[[(2-thienyl)hydroxyphosphinyl)methyl]pentanedioic
- acid;
- 2-[[[(3-thienyl)hydroxyphosphinyl)methyl]pentanedioic
- acid;
- 2-[[[(4-thienyl)hydroxyphosphinyl)methyl]pentanedioic
- 25 acid; and
- pharmaceutically acceptable salts and hydrates thereof.

55. The method of claim 46, wherein R_1 is 1-naphthyl, 2-naphthyl, or C_1 - C_4 straight or branched chain alkyl substituted with 1-naphthyl or 2-naphthyl.

- 5 56. The method of claim 55, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 2 - [[(1-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 10 2 - [[(2-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2 - [[(1-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2 - [[(2-naphthyl)methylhydroxyphosphinyl]methyl]pentanedioic acid;
- 15 2 - [[(1-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2 - [[(2-naphthyl)ethylhydroxyphosphinyl]methyl]pentanedioic acid;
- 20 2 - [[(1-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2 - [[(2-naphthyl)propylhydroxyphosphinyl]methyl]pentanedioic acid;
- 2 - [[(1-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic acid;
- 25 2 - [[(2-naphthyl)butylhydroxyphosphinyl]methyl]pentanedioic acid;

210

dioic acid; and

pharmaceutically acceptable salts and hydrates thereof.

57. The method of claim 45, wherein:

5 R_2 is selected from the group consisting of
hydrogen, C_1 - C_9 straight or branched chain alkyl, C_2 - C_9
straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_9
cycloalkenyl, benzyl and phenyl, wherein said R_2 is
unsubstituted or substituted with C_3 - C_8 cycloalkyl, C_5 - C_9
10 cycloalkenyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6
straight or branched chain alkenyl, C_1 - C_4 alkoxy, phenyl
or mixtures thereof.

58. The method of claim 57, wherein:

15 R_1 is hydrogen, C_1 - C_4 straight or branched chain
alkyl, C_2 - C_4 straight or branched chain alkenyl, C_3 - C_8
cycloalkyl, C_5 - C_9 cycloalkenyl, benzyl or phenyl, wherein
said R_1 is unsubstituted or substituted with carboxy, C_3 -
 C_8 cycloalkyl, C_5 - C_9 cycloalkenyl, halo, hydroxy, nitro,
20 trifluoromethyl, C_1 - C_6 straight or branched chain alkyl,
 C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 -
 C_4 alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl
or mixtures thereof.

25 59. The method of claim 58, wherein the glutamate-
derived hydroxyphosphinyl derivative is selected from the

SUBSTITUTE SHEET (RULE 26)

group consisting of:

- 3-(methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(propylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 5 3-(butylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(cyclohexylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((cyclohexyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(phenylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 10 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(phenylethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(phenylpropylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(phenylbutylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl)-2-
- 15 phenylpropanoic acid;
- 3-(phenylprop-2-enylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-propylpropanoic acid;
- 20 3-(benzylhydroxyphosphinyl)-2-butylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-cyclohexylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(cyclohexyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 25 3-(benzylhydroxyphosphinyl)-2-benzylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylethylpropanoic acid;

212

- 3-(benzylhydroxyphosphinyl)-2-phenylpropylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylbutylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(2,3,4-trimethoxyphenyl)-
propanoic acid;
5 3-(benzylhydroxyphosphinyl)-2-phenylprop-2-enylpropanoic
acid; and
pharmaceutically acceptable salts and hydrates thereof.

60. The method of claim 45, wherein at least one of
10 R₁ and R₂ is 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-
furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl,
3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or C₁-C₄
straight or branched chain alkyl substituted with 2-
indolyl 3-indolyl, 4-indolyl, 2-furyl, 3-furyl,
15 tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-
pyridyl or 4-pyridyl.

61. The method of claim 60, wherein the glutamate-
derived hydroxyphosphinyl derivative is selected from the
20 group consisting of:
3-[(2-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
3-[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;
25 3-[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic
acid;

SUBSTITUTE SHEET (RULE 26)

213

- 3-[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 5 3-[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 10 3-[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 15 3-[(4-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 20 3-[(2-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 25 3-[(4-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

SUBSTITUTE SHEET (RULE 26)

214

- 3-[(3-thienyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-thienyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 5 3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)methylpropanoic
- 10 acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propylpropanoic acid;
- 15 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)ethylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(tetrahydrofuranyl)propyl
- 20 propanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(2-indolyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-indolyl)methylpropanoic acid;
- 25 3-(benzylhydroxyphosphinyl)-2-(4-indolyl)methylpropanoic acid;

SUBSTITUTE SHEET (RULE 26)

215

- 3 - (benzylhydroxyphosphinyl) - 2 - (3-indolyl) ethylpropanoic acid;
- 3 - (benzylhydroxyphosphinyl) - 2 - (3-indolyl) propylpropanoic acid;
- 5 3 - (benzylhydroxyphosphinyl) - 2 - (2-thienyl) methylpropanoic acid;
- 3 - (benzylhydroxyphosphinyl) - 2 - (3-thienyl) methylpropanoic acid;
- 3 - (benzylhydroxyphosphinyl) - 2 - (4-thienyl) methylpropanoic
- 10 acid;
- 3 - (benzylhydroxyphosphinyl) - 2 - (3-thienyl) ethylpropanoic acid;
- 3 - (benzylhydroxyphosphinyl) - 2 - (3-thienyl) propylpropanoic acid; and
- 15 pharmaceutically acceptable salts and hydrates thereof.

62. The method of claim 57, wherein R₁ is 1-naphthyl, 2-naphthyl, or C₁-C₄ straight or branched chain alkyl substituted with 1-naphthyl or 2-naphthyl.

20

63. The method of claim 62, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 3 - ((1-naphthyl)hydroxyphosphinyl) - 2-phenylpropanoic acid;
- 25 3 - ((2-naphthyl)hydroxyphosphinyl) - 2-phenylpropanoic acid;
- 3 - ((1-naphthyl)methylhydroxyphosphinyl) - 2-phenylpropanoic

SUBSTITUTE SHEET (RULE 26)

216

- acid;
3 - ((2-naphthyl)methylhydroxyphosphinyl) - 2-phenylpropanoic
acid;
3 - ((1-naphthyl)ethylhydroxyphosphinyl) - 2-phenylpropanoic
5 acid;
3 - ((2-naphthyl)ethylhydroxyphosphinyl) - 2-phenylpropanoic
acid;
3 - ((1-naphthyl)propylhydroxyphosphinyl) - 2-phenylpropanoic
acid;
10 3 - ((2-naphthyl)propylhydroxyphosphinyl) - 2-phenylpropanoic
acid;
3 - ((1-naphthyl)butylhydroxyphosphinyl) - 2-phenylpropanoic
acid;
3 - ((2-naphthyl)butylhydroxyphosphinyl) - 2-phenylpropanoic
15 acid; and
pharmaceutically acceptable salts and hydrates thereof.

64. The method of claim 44, wherein X is O.

- 20 65. The method of claim 64, wherein R₂ is
substituted with carboxy.

66. The method of claim 65, wherein the glutamate-
derived hydroxyphosphinyl derivative is selected from the
25 group consisting of:

2 - [[methylhydroxyphosphinyl]oxy]pentanedioic acid;

217

- 2- [[ethylhydroxyphosphinyl]oxy]pentanedioic acid;
2- [[propylhydroxyphosphinyl]oxy]pentanedioic acid;
2- [[butylhydroxyphosphinyl]oxy]pentanedioic acid;
2- [[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;
5 2- [[(cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2- [[phenylhydroxyphosphinyl]oxy]pentanedioic acid;
2- [[benzylhydroxyphosphinyl]oxy]pentanedioic acid;
2- [[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;
10 2- [[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;
2- [[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;
2- [[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic
acid;
2- [[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic
15 acid;
2- [[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic
acid;
2- [[(pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic
acid;
20 2- [[(methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic
acid;
2- [[(2,3,4-trimethoxyphenyl)hydroxyphosphinyl]oxy]-
pentanedioic acid;
2- [[(1-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
25 2- [[(2-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2- [[(1-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic

SUBSTITUTE SHEET (RULE 26)

218

- acid;
2- [[(2-naphthyl) methylhydroxyphosphinyl] oxy] pentanedioic
acid;
2- [[(1-naphthyl) ethylhydroxyphosphinyl] oxy] pentanedioic
5 acid;
2- [[(2-naphthyl) ethylhydroxyphosphinyl] oxy] pentanedioic
acid;
2- [[(1-naphthyl) propylhydroxyphosphinyl] oxy] pentanedioic
acid;
10 2- [[(2-naphthyl) propylhydroxyphosphinyl] oxy] pentanedioic
acid;
2- [[(1-naphthyl) butylhydroxyphosphinyl] oxy] pentanedioic
acid;
2- [[(2-naphthyl) butylhydroxyphosphinyl] oxy] pentanedioic
15 acid;
2- [[(phenylprop-2-enyl) hydroxyphosphinyl] oxy] pentanedioic
acid;
2- [[(benzylhydroxyphosphinyl] oxy] pentanedioic acid;
2- [[((hydroxy) phenylmethyl) hydroxyphosphinyl] oxy] pentane-
20 dioic acid;
2- [[(3-methylbenzyl) hydroxyphosphinyl] oxy] pentanedioic
acid;
2- [[(4-fluorophenyl) hydroxyphosphinyl] oxy] pentanedioic
acid;
25 2- [[(2-fluorobenzyl) hydroxyphosphinyl] oxy] pentanedioic
acid;

SUBSTITUTE SHEET (RULE 26)

219

- 2-(phosphono)oxy]pentanedioic acid;
2-[[(3-trifluoromethylbenzyl) hydroxyphosphinyl]oxy] -
pentanedioic acid;
2-[[methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
5 2-[[ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[butylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[cyclohexylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
10 2-[[(cyclohexyl) methylhydroxyphosphinyl]oxy] -2-phenyl-
ethanoic acid;
2-[[phenylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[phenylethylhydroxyphosphinyl]oxy]-2-phenylethanoic
15 acid;
2-[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
2-[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic
acid;
20 2-[[(2,3,4-trimethoxyphenyl) -3-hydroxyphosphinyl]oxy] -2-
phenylethanoic acid;
2-[[(1-naphthyl) hydroxyphosphinyl]oxy] -2-phenylethanoic
acid;
2-[[(2-naphthyl) hydroxyphosphinyl]oxy] -2-phenylethanoic
25 acid;
2-[[(1-naphthyl) methylhydroxyphosphinyl]oxy] -2-phenyl-

SUBSTITUTE SHEET (RULE 26)

- ethanoic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 5 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 10 2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 15 2-[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;
- 20 2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;
- 25 2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]decanedioic acid;

221

- 2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-methylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-ethylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-propylethanoic acid;
5 2-[[benzylhydroxyphosphinyl]oxy]-2-butylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-cyclohexylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(cyclohexyl)methyl-
ethanoic acid;
10 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-benzylethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylpropylethanoic
15 acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-phenylbutylethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2,3,4-trimethoxy-
phenyl)ethanoic acid;
20 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethanoic
acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)methyl-
25 ethanoic acid;
2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)methyl-

SUBSTITUTE SHEET (RULE 26)

222

- ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethyl-ethanoic acid;
- 5 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propyl-ethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enyl-ethanoic acid;
- 15 2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 20 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 25 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-

SUBSTITUTE SHEET (RULE 26)

223

- pentanedioic acid;
2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-
pentanedioic acid;
2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-
5 pentanedioic acid;
2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
10 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic
15 acid;
2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
20 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic
25 acid; and
pharmaceutically acceptable salts and hydrates thereof.

SUBSTITUTE SHEET (RULE 26)

67. The method of claim 64, wherein:

R₂ is selected from the group consisting of hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl and phenyl, wherein said R₂ is unsubstituted or substituted with C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl, C₁-C₄ alkoxy, phenyl or mixtures thereof.

10

68. The method of claim 67, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 15 2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 20 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 25 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-

225

- phenylethanoic acid;
- 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-
- 5 ethanoic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 10 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-
- 15 ethanoic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 20 2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)methyl-
- 25 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)methyl-

SUBSTITUTE SHEET (RULE 26)

- ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethyl-
- 5 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-methylethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-indolyl)methyl-
- 15 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-indolyl)methyl-ethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-indolyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-thienyl)methyl-
- 25 ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)methyl-

ethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(4-thienyl)methyl-ethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)ethyl-

5 ethanoic acid;

2-[[benzylhydroxyphosphinyl]oxy]-2-(3-thienyl)propyl-ethanoic acid; and

pharmaceutically acceptable salts and hydrates thereof.

10 69. The method of claim 44, wherein X is NR₁.

70. The method of claim 69, wherein R₂ is substituted with carboxy.

15 71. The method of claim 70, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;

20 2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[cyclohexyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

25 2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;

2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;

- 2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;
2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic
acid;
2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;
5 2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic
10 acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentane-
dioic acid;
2-[[[(methoxybenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;
15 2-[[[(2,3,4-trimethoxyphenyl)hydroxyphosphinyl]amino]-
pentanedioic acid;
2-[[[(1-naphthyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]amino]pentanedioic
20 acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]amino]pentane-
dioic acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]amino]pentane-
dioic acid;
25 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]amino]pentanedioic
acid;

- 2- [[(2-naphthyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(1-naphthyl) propylhydroxyphosphinyl] amino] pentanedioic acid;
- 5 2- [[(2-naphthyl) propylhydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(1-naphthyl) butylhydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(2-naphthyl) butylhydroxyphosphinyl] amino] pentanedioic acid;
- 10 acid;
- 2- [[(phenylprop-2-enyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(benzylhydroxyphosphinyl) amino] pentanedioic acid;
- 2- [[(2-fluorobenzyl) hydroxyphosphinyl] amino] -2-pentanedioic acid;
- 15 acid;
- 2- [[((hydroxy) phenylmethyl) hydroxyphosphinyl] amino] -pentanedioic acid;
- 2- [[(3-methylbenzyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 20 2- [[(4-fluorophenyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2- [(phosphono) amino] pentanedioic acid;
- 2- [[(3-trifluoromethylbenzyl) hydroxyphosphinyl] amino] -pentanedioic acid;
- 25 2- [(methylhydroxyphosphinyl) amino] hexanedioic acid;
- 2- [(benzylhydroxyphosphinyl) amino] hexanedioic acid;

230

- 2- [(methylhydroxyphosphinyl) amino] heptanedioic acid;
2- [(benzylhydroxyphosphinyl) amino] heptanedioic acid;
2- [(methylhydroxyphosphinyl) amino] octanedioic acid;
2- [(benzylhydroxyphosphinyl) amino] octanedioic acid;
5 2- [(methylhydroxyphosphinyl) amino] nonanedioic acid;
2- [(benzylhydroxyphosphinyl) amino] nonanedioic acid;
2- [(methylhydroxyphosphinyl) amino] decanedioic acid;
2- [(benzylhydroxyphosphinyl) amino] decanedioic acid;
3- [[(2-pyridyl) methylhydroxyphosphinyl] amino] pentanedioic
10 acid;
3- [[(3-pyridyl) methylhydroxyphosphinyl] amino] pentanedioic
acid;
3- [[(4-pyridyl) methylhydroxyphosphinyl] amino] pentanedioic
acid;
15 3- [[(3-pyridyl) ethylhydroxyphosphinyl] amino] pentanedioic
acid;
3- [[(3-pyridyl) propylhydroxyphosphinyl] amino] pentanedioic
acid;
3- [[(tetrahydrofuranyl) methylhydroxyphosphinyl] amino] -
20 pentanedioic acid;
3- [[(tetrahydrofuranyl) ethylhydroxyphosphinyl] amino] -
pentanedioic acid;
3- [[(tetrahydrofuranyl) propylhydroxyphosphinyl] amino] -
pentanedioic acid;
25 3- [[(2-indolyl) methylhydroxyphosphinyl] amino] pentanedioic
acid;

SUBSTITUTE SHEET (RULE 26)

231

3-[[(3-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[(4-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

5 3-[[(3-indolyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[(3-indolyl)propylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[(2-thienyl)methylhydroxyphosphinyl]amino]pentanedioic
10 acid;

3-[[(3-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[(4-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

15 3-[[(3-thienyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;

3-[[(3-thienyl)propylhydroxyphosphinyl]amino]pentanedioic acid; and

pharmaceutically acceptable salts and hydrates thereof.

20

72. A method of effecting a neuronal activity in an animal, comprising administering an effective amount of a NAALADase inhibitor to said animal.

25 73. The method of claim 72, wherein the neuronal activity is selected from the group consisting of

stimulation of damaged neurons, promotion of neuronal regeneration, prevention of neurodegeneration, and treatment of a neurological disorder.

5 74. The method of claim 73, wherein the neurological disorder is selected from the group consisting of peripheral neuropathy caused by physical injury or disease state, traumatic brain injury, physical damage to the spinal cord, stroke associated with brain
10 damage, demyelinating disease and neurological disorder relating to neurodegeneration.

75. The method of claim 74, wherein the peripheral neuropathy is caused by Guillain-Barré syndrome.

15

76. The method of claim 74, wherein the demyelinating disease is multiple sclerosis.

77. The method of claim 74, wherein the
20 neurological disorder relating to neurodegeneration is selected from the group consisting of Alzheimer's Disease, Parkinson's Disease, and amyotrophic lateral sclerosis.

25 78. The method of claim 72, wherein the NAALADase inhibitor is administered in combination with at least


233

one additional therapeutic agent.

79. The method of claim 72, wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative, an acidic peptide analog or a mixture thereof.

80. The method of claim 79, wherein the NAALADase inhibitor is an acidic peptide analog selected from the group consisting of Asp-Glu, Glu-Glu, Gly-Glu, gamma-Glu-Glu and Glu-Glu-Glu.

81. The method of claim 79, wherein the NAALADase inhibitor is a glutamate-derived hydroxyphosphinyl derivative of formula I:



I

20

or a pharmaceutically acceptable salt or hydrate thereof, wherein:

R₁ is selected from the group consisting of hydrogen, C₁-C₈ straight or branched chain alkyl, C₂-C₈ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl and Ar, wherein said R₁ is unsubstituted or

SUBSTITUTE SHEET (RULE 26)

substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, 5 phenoxy, benzyloxy, amino, Ar or mixtures thereof;

X is CR₃R₄, O or NR₁;

R₃ and R₄ are independently selected from the group consisting of hydrogen, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, Ar, halo and mixtures 10 thereof;

R₂ is selected from the group consisting of hydrogen, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl and Ar, wherein said R₂ is unsubstituted or 15 substituted with carboxy, C₃-C₈ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, 20 phenoxy, benzyloxy, amino, Ar or mixtures thereof;

Ar is selected from the group consisting of 1-naphthyl, 2-naphthyl, 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 25 benzyl and phenyl, wherein said Ar is unsubstituted or substituted with halo, hydroxy, nitro, trifluoromethyl,

235

C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₆ alkoxy, C₂-C₆ alkenyloxy, phenoxy, benzyloxy, amino or mixtures thereof.

5 82. The method of claim 81, wherein X is CH₂.

83. The method of claim 82, wherein R₂ is substituted with carboxy.

10 84. The method of claim 83, wherein:

R₁ is hydrogen, C₁-C₄ straight or branched chain alkyl, C₂-C₄ straight or branched chain alkenyl, C₃-C₆ cycloalkyl, C₅-C₇ cycloalkenyl, benzyl or phenyl, wherein said R₁ is unsubstituted or substituted with carboxy, C₃-
15 C₆ cycloalkyl, C₅-C₇ cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof; and

20 R₂ is C₁-C₂ alkyl.

85. The method of claim 84, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 25 2-(phosphonomethyl)pentanedioic acid;
2-(phosphonomethyl)succinic acid;

SUBSTITUTE SHEET (RULE 26)

236

- 2-[[(2-carboxyethyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[methylhydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[ethylhydroxyphosphinyl] methyl] pentanedioic acid;
- 5 2-[[propylhydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[butylhydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(cyclohexyl) methylhydroxyphosphinyl] methyl] pentanedioic acid;
- 10 2-[[phenylhydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(benzylhydroxyphosphinyl) methyl] pentanedioic acid;
- 2-[[(phenylmethyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(phenylethyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 15 2-[[(phenylpropyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(phenylbutyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 20 2-[[(4-methylbenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(4-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 2-[[(2-fluorobenzyl) hydroxyphosphinyl] methyl] pentanedioic acid;
- 25 2-[[(pentafluorobenzyl) hydroxyphosphinyl] methyl] pentane-

SUBSTITUTE SHEET (RULE 26)

237

- dioic acid;
- 2-[[[(methoxybenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2,3,4-trimethoxyphenyl)hydroxyphosphinyl)methyl]-
- 5 pentanedioic acid;
- 2-[[[(phenylprop-2-enyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 10 2-[[[(hydroxy)phenylmethyl)hydroxyphosphinyl)methyl]-pentanedioic acid;
- 2-[[[(3-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic acid;
- 2-[[[(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic
- 15 acid;
- 2-[[[(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl]-pentanedioic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

- 20 86. The method of claim 85, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 2-(phosphonomethyl)pentanedioic acid;
- 2-(phosphonomethyl)succinic acid;
- 25 2-[[[(2-carboxyethyl)hydroxyphosphinyl)methyl]pentanedioic acid;

SUBSTITUTE SHEET (RULE 26)

- 2 - [(benzylhydroxyphosphinyl)methyl]pentanedioic acid;
2 - [(phenylhydroxyphosphinyl)methyl]pentanedioic acid;
2 - [((hydroxy)phenylmethyl)hydroxyphosphinyl)methyl] -
pentanedioic acid;
- 5 2 - [(butylhydroxyphosphinyl)methyl]pentanedioic acid;
2 - [(3-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
2 - [(3-phenylpropylhydroxyphosphinyl)methyl]pentanedioic
acid;
- 10 2 - [(4-fluorophenyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
2 - [(methylhydroxyphosphinyl)methyl]pentanedioic acid;
2 - [(phenylethylhydroxyphosphinyl)methyl]pentanedioic
acid;
- 15 2 - [(4-methylbenzyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
2 - [(4-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
2 - [(4-methoxybenzyl)hydroxyphosphinyl)methyl]pentane-
- 20 dioic acid;
2 - [(3-trifluoromethylbenzyl)hydroxyphosphinyl)methyl] -
pentanedioic acid;
2 - [(2-fluorobenzyl)hydroxyphosphinyl)methyl]pentanedioic
acid;
- 25 2 - [(pentafluorobenzyl)hydroxyphosphinyl)methyl]pentane-
dioic acid;

239

2-[[(phenylprop-2-enyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(aminomethyl)hydroxyphosphinyl]methyl]pentanedioic acid;

5 2-[[(aminoethyl)hydroxyphosphinyl]methyl]pentanedioic acid;

2-[[(aminopropyl)hydroxyphosphinyl]methyl]pentanedioic acid; and

pharmaceutically acceptable salts and hydrates thereof.

10

87. The method of claim 86, wherein the glutamate-derived hydroxyphosphinyl derivative is 2-(phosphonomethyl)pentanedioic acid or a pharmaceutically acceptable salt or hydrate thereof.

15

88. The method of claim 83, wherein:

R_1 is hydrogen, C_1 - C_4 straight or branched chain alkyl, C_2 - C_4 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, benzyl or phenyl, wherein
20 said R_1 is unsubstituted or substituted with carboxy, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 - C_4 alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl
25 or mixtures thereof; and

R_2 is C_3 - C_9 alkyl.

89. The method of claim 88, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 2-[(methylhydroxyphosphinyl)methyl]hexanedioic acid;
- 5 2-[(benzylhydroxyphosphinyl)methyl]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]octanedioic acid;
- 10 2-[(methylhydroxyphosphinyl)methyl]nonanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]nonanedioic acid;
- 2-[(methylhydroxyphosphinyl)methyl]decanedioic acid;
- 2-[(benzylhydroxyphosphinyl)methyl]decanedioic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

15

90. The method of claim 83, wherein R_1 is 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl or C_1 - C_4 straight
- 20 or branched chain alkyl substituted with 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl or 4-pyridyl.

91. The method of claim 90, wherein the glutamate-
- 25 derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 2-[[[(2-pyridyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 5 2-[[[(4-pyridyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]methyl]pentane-
10 dioic acid;
- 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]methyl]-
pentanedioic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]methyl]-
pentanedioic acid;
- 15 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]methyl]-
pentanedioic acid;
- 2-[[[(2-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
pentanedioic acid;
- 2-[[[(3-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
20 pentanedioic acid;
- 2-[[[(4-tetrahydropyranyl)hydroxyphosphinyl]methyl]-
pentanedioic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 25 2-[[[(3-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;

- 2-[[[(4-indolyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 5 2-[[[(3-indolyl)propylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(2-thienyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]methyl]pentane-
10 dioic acid;
- 2-[[[(4-thienyl)methylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 15 2-[[[(3-thienyl)propylhydroxyphosphinyl]methyl]pentane-
dioic acid;
- 2-[[[(2-pyridyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
- 2-[[[(3-pyridyl)hydroxyphosphinyl]methyl]pentanedioic
20 acid;
- 2-[[[(4-pyridyl)hydroxyphosphinyl]methyl]pentanedioic
acid;
- 2-[[[(tetrahydrofuranyl)hydroxyphosphinyl]methyl]pentane-
dioic acid;
- 25 2-[[[(2-indolyl)hydroxyphosphinyl]methyl]pentanedioic
acid;

243

- 2-[[[(3-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(4-indolyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 5 2-[[[(2-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(3-thienyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(4-thienyl)hydroxyphosphinyl]methyl]pentanedioic
- 10 acid; and
- pharmaceutically acceptable salts and hydrates thereof.

92. The method of claim 83, wherein R₁ is 1-naphthyl, 2-naphthyl, or C₁-C₄ straight or branched chain

15 alkyl substituted with 1-naphthyl or 2-naphthyl.

93. The method of claim 92, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 20 2-[[[(1-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(2-naphthyl)hydroxyphosphinyl]methyl]pentanedioic acid;
- 2-[[[(1-naphthyl)methylhydroxyphosphinyl]methyl]pentane-
- 25 dioic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]methyl]pentane-

244

- dioic acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]methyl]pentane-
dioic acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]methyl]pentane-
5 dioic acid;
2-[[[(1-naphthyl)propylhydroxyphosphinyl]methyl]pentane-
dioic acid;
2-[[[(2-naphthyl)propylhydroxyphosphinyl]methyl]pentane-
dioic acid;
10 2-[[[(1-naphthyl)butylhydroxyphosphinyl]methyl]pentane-
dioic acid;
2-[[[(2-naphthyl)butylhydroxyphosphinyl]methyl]pentane-
dioic acid; and
pharmaceutically acceptable salts and hydrates thereof.
15

94. The method of claim 82, wherein:

R₂ is selected from the group consisting of
hydrogen, C₁-C₉ straight or branched chain alkyl, C₂-C₉
straight or branched chain alkenyl, C₃-C₈ cycloalkyl, C₅-C₇
20 cycloalkenyl, benzyl and phenyl, wherein said R₂ is
unsubstituted or substituted with C₃-C₈ cycloalkyl, C₅-C₇
cycloalkenyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆
straight or branched chain alkenyl, C₁-C₄ alkoxy, phenyl
or mixtures thereof.

25

95. The method of claim 94, wherein:

245

R_1 is hydrogen, C_1 - C_4 straight or branched chain alkyl, C_2 - C_4 straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, benzyl or phenyl, wherein said R_1 is unsubstituted or substituted with carboxy, C_3 -
 5 C_8 cycloalkyl, C_5 - C_7 cycloalkenyl, halo, hydroxy, nitro, trifluoromethyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl, C_1 - C_4 alkoxy, C_2 - C_4 alkenyloxy, phenoxy, benzyloxy, amino, benzyl, phenyl or mixtures thereof.

10

96. The method of claim 95, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 3-(methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 15 3-(ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(propylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(butylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(cyclohexylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((cyclohexyl)methylhydroxyphosphinyl)-2-phenylpropanoic
 20 acid;
- 3-(phenylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(phenylethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-(phenylpropylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 25 3-(phenylbutylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl)-2-

SUBSTITUTE SHEET (RULE 26)

246

- phenylpropanoic acid;
3-(phenylprop-2-enylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-ethylpropanoic acid;
5 3-(benzylhydroxyphosphinyl)-2-propylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-butylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-cyclohexylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-(cyclohexyl)methylpropanoic acid;
10 3-(benzylhydroxyphosphinyl)-2-phenylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-benzylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylethylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylpropylpropanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylbutylpropanoic acid;
15 3-(benzylhydroxyphosphinyl)-2-(2,3,4-trimethoxyphenyl)-propanoic acid;
3-(benzylhydroxyphosphinyl)-2-phenylprop-2-enylpropanoic acid; and
pharmaceutically acceptable salts and hydrates thereof.

20

97. The method of claim 82, wherein at least one of R_1 and R_2 is 2-indolyl, 3-indolyl, 4-indolyl, 2-furyl, 3-furyl, tetrahydrofuranyl, tetrahydropyranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or C_1 - C_4
25 straight or branched chain alkyl substituted with 2-indolyl 3-indolyl, 4-indolyl, 2-furyl, 3-furyl,

SUBSTITUTE SHEET (RULE 26)

247

tetrahydrofuranyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl or 4-pyridyl.

98. The method of claim 97, wherein the glutamate-
5 derived hydroxyphosphinyl derivative is selected from the group consisting of:

3-[(2-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic
10 acid;

3-[(4-pyridyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(3-pyridyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;

15 3-[(3-pyridyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(tetrahydrofuranyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(tetrahydrofuranyl)ethylhydroxyphosphinyl]-2-phenyl
20 propanoic acid;

3-[(tetrahydrofuranyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;

3-[(2-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

25 3-[(3-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;

- 3-[(4-indolyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-indolyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 5 3-[(3-indolyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(2-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic
10 acid;
- 3-[(4-thienyl)methylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-[(3-thienyl)ethylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 15 3-[(3-thienyl)propylhydroxyphosphinyl]-2-phenylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(2-pyridyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)methylpropanoic
20 acid;
- 3-(benzylhydroxyphosphinyl)-2-(4-pyridyl)methylpropanoic acid;
- 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)ethylpropanoic acid;
- 25 3-(benzylhydroxyphosphinyl)-2-(3-pyridyl)propylpropanoic acid;

249

- 3 - (benzylhydroxyphosphinyl) -2 - (tetrahydrofuranyl) methyl
propanoic acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (tetrahydrofuranyl) ethyl
propanoic acid;
- 5 3 - (benzylhydroxyphosphinyl) -2 - (tetrahydrofuranyl) propyl
propanoic acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (2-indolyl) methylpropanoic
acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (3-indolyl) methylpropanoic
10 acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (4-indolyl) methylpropanoic
acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (3-indolyl) ethylpropanoic
acid;
- 15 3 - (benzylhydroxyphosphinyl) -2 - (3-indolyl) propylpropanoic
acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (2-thienyl) methylpropanoic
acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (3-thienyl) methylpropanoic
20 acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (4-thienyl) methylpropanoic
acid;
- 3 - (benzylhydroxyphosphinyl) -2 - (3-thienyl) ethylpropanoic
acid;
- 25 3 - (benzylhydroxyphosphinyl) -2 - (3-thienyl) propylpropanoic
acid; and

SUBSTITUTE SHEET (RULE 26)

pharmaceutically acceptable salts and hydrates thereof.

99. The method of claim 94, wherein R_1 is 1-naphthyl, 2-naphthyl, or C_1 - C_4 straight or branched chain alkyl substituted with 1-naphthyl or 2-naphthyl.

100. The method of claim 99, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 10 3-((1-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)hydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((1-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)methylhydroxyphosphinyl)-2-phenylpropanoic
- 15 acid;
- 3-((1-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)ethylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 20 3-((1-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((2-naphthyl)propylhydroxyphosphinyl)-2-phenylpropanoic acid;
- 3-((1-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic
- 25 acid;
- 3-((2-naphthyl)butylhydroxyphosphinyl)-2-phenylpropanoic

251

acid; and

pharmaceutically acceptable salts and hydrates thereof.

101. The method of claim 81, wherein X is O.

5

102. The method of claim 101, wherein R₂ is substituted with carboxy.

103. The method of claim 102, wherein the glutamate-
10 derived hydroxyphosphinyl derivative is selected from the group consisting of:

- 2-[[methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 15 2-[[butylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[cyclohexylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(cyclohexyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[phenylhydroxyphosphinyl]oxy]pentanedioic acid;
- 20 2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[phenylethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[phenylpropylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[phenylbutylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[[(4-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic
- 25 acid;
- 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic

- acid;
2-[[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(pentafluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic
5 acid;
2-[[[(methoxybenzyl)hydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(2,3,4-trimethoxyphenyl)hydroxyphosphinyl]oxy]-
pentanedioic acid;
10 2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]pentanedioic acid;
2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]pentanedioic
15 acid;
2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
20 2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic
25 acid;
2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]pentanedioic

- acid;
- 2-[[(phenylprop-2-enyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]pentanedioic acid;
- 5 2-[[[(hydroxy)phenylmethyl]hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(3-methylbenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(4-fluorophenyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 10 2-[[(2-fluorobenzyl)hydroxyphosphinyl]oxy]pentanedioic acid;
- 2- (phosphono)oxy]pentanedioic acid;
- 2-[[(3-trifluoromethylbenzyl)hydroxyphosphinyl]oxy] -
- 15 pentanedioic acid;
- 2-[[methylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 2-[[ethylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 2-[[propylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 2-[[butylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 20 2-[[cyclohexylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 2-[[(cyclohexyl)methylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 2-[[phenylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]oxy] -2-phenylethanoic acid;
- 2-[[phenylethylhydroxyphosphinyl]oxy] -2-phenylethanoic

- acid;
- 2-[[phenylpropylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[phenylbutylhydroxyphosphinyl]oxy]-2-phenylethanoic
- 5 acid;
- 2-[[[(2,3,4-trimethoxyphenyl)-3-hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 10 2-[[[(2-naphthyl)hydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(1-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(2-naphthyl)methylhydroxyphosphinyl]oxy]-2-phenyl-
- 15 ethanoic acid;
- 2-[[[(1-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(2-naphthyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 20 2-[[[(1-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(2-naphthyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(1-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenyl-
- 25 ethanoic acid;
- 2-[[[(2-naphthyl)butylhydroxyphosphinyl]oxy]-2-phenyl-

- ethanoic acid;
- 2-[[phenylprop-2-enylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[(methylhydroxyphosphinyl)oxy]hexanedioic acid;
- 5 2-[(benzylhydroxyphosphinyl)oxy]hexanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]heptanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]heptanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]octanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]octanedioic acid;
- 10 2-[(methylhydroxyphosphinyl)oxy]nonanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]nonanedioic acid;
- 2-[(methylhydroxyphosphinyl)oxy]decanedioic acid;
- 2-[(benzylhydroxyphosphinyl)oxy]decanedioic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-methylethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]oxy]-2-ethylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-propylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-butylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-cyclohexylethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]oxy]-2-(cyclohexyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-benzylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylethylethanoic
- 25 acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylpropylethanoic

- acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylbutylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2,3,4-trimethoxy-phenyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)propyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(1-naphthyl)butyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-naphthyl)butyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-phenylprop-2-enyl-

257

- ethanoic acid;
- 2-[[(2-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(3-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 5 2-[[(4-pyridyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 10 2-[[(3-pyridyl)propylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy] - pentanedioic acid;
- 2-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy] - pentanedioic acid;
- 15 2-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy] - pentanedioic acid;
- 2-[[(2-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 20 2-[[(3-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(4-indolyl)methylhydroxyphosphinyl]oxy]pentanedioic acid;
- 2-[[(3-indolyl)ethylhydroxyphosphinyl]oxy]pentanedioic acid;
- 25 2-[[(3-indolyl)propylhydroxyphosphinyl]oxy]pentanedioic

SUBSTITUTE SHEET (RULE 26)

- acid;
2-[[(2-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[(3-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic
5 acid;
2-[[(4-thienyl)methylhydroxyphosphinyl]oxy]pentanedioic
acid;
2-[[(3-thienyl)ethylhydroxyphosphinyl]oxy]pentanedioic
acid;
10 2-[[(3-thienyl)propylhydroxyphosphinyl]oxy]pentanedioic
acid; and
pharmaceutically acceptable salts and hydrates thereof.

104. The method of claim 101, wherein:

- 15 R_2 is selected from the group consisting of
hydrogen, C_1 - C_8 straight or branched chain alkyl, C_2 - C_8
straight or branched chain alkenyl, C_3 - C_8 cycloalkyl, C_5 - C_7
cycloalkenyl, benzyl and phenyl, wherein said R_2 is
unsubstituted or substituted with C_1 - C_8 cycloalkyl, C_5 - C_7
20 cycloalkenyl, C_1 - C_8 straight or branched chain alkyl, C_2 - C_8
straight or branched chain alkenyl, C_1 - C_4 alkoxy, phenyl
or mixtures thereof.

105. The method of claim 104, wherein the glutamate-
25 derived hydroxyphosphinyl derivative is selected from the
group consisting of:

- 2-[[[(2-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 5 2-[[[(4-pyridyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-pyridyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-pyridyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 10 2-[[[(tetrahydrofuranyl)methylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 15 2-[[[(tetrahydrofuranyl)propylhydroxyphosphinyl]oxy]-2-phenylethanoic acid;
- 2-[[[(2-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 20 2-[[[(4-indolyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-indolyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 25 2-[[[(3-indolyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;

260

- 2-[[[(2-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 5 2-[[[(4-thienyl)methylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-thienyl)ethylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 2-[[[(3-thienyl)propylhydroxyphosphinyl]oxy]-2-phenyl-ethanoic acid;
- 10 2-[[benzylhydroxyphosphinyl]oxy]-2-(2-pyridyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)methyl-ethanoic acid;
- 15 2-[[benzylhydroxyphosphinyl]oxy]-2-(4-pyridyl)methyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)ethyl-ethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(3-pyridyl)propyl-ethanoic acid;
- 20 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-methylethanoic acid;
- 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-ethylethanoic acid;
- 25 2-[[benzylhydroxyphosphinyl]oxy]-2-(tetrahydrofuranyl)-propylethanoic acid;

SUBSTITUTE SHEET (RULE 26)

261

- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (2-indolyl) methyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-indolyl) methyl-ethanoic acid;
- 5 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (4-indolyl) methyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-indolyl) ethyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-indolyl) propyl-
- 10 ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (2-thienyl) methyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-thienyl) methyl-ethanoic acid;
- 15 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (4-thienyl) methyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-thienyl) ethyl-ethanoic acid;
- 2 - [[benzylhydroxyphosphinyl]oxy] - 2 - (3-thienyl) propyl-
- 20 ethanoic acid; and
- pharmaceutically acceptable salts and hydrates thereof.

106. The method of claim 81, wherein X is NR_1 .

25

107. The method of claim 106, wherein R_2 is

262

substituted with carboxy.

108. The method of claim 107, wherein the glutamate-derived hydroxyphosphinyl derivative is selected from the group consisting of:
- 2-[[methylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[ethylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[propylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[butylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[cyclohexylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[[(cyclohexyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[phenylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[benzylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[phenylethylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[phenylpropylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[phenylbutylhydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[[(4-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[[(4-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[[(2-fluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;
 - 2-[[[(pentafluorobenzyl)hydroxyphosphinyl]amino]pentanedioic acid;

263

- 2- [[(methoxybenzyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(2,3,4-trimethoxyphenyl) hydroxyphosphinyl] amino] - pentanedioic acid;
- 5 2- [[(1-naphthyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(2-naphthyl) hydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(1-naphthyl) methylhydroxyphosphinyl] amino] pentane-
- 10 dioic acid;
- 2- [[(2-naphthyl) methylhydroxyphosphinyl] amino] pentane-dioic acid;
- 2- [[(1-naphthyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 15 2- [[(2-naphthyl) ethylhydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(1-naphthyl) propylhydroxyphosphinyl] amino] pentane-dioic acid;
- 2- [[(2-naphthyl) propylhydroxyphosphinyl] amino] pentane-
- 20 dioic acid;
- 2- [[(1-naphthyl) butylhydroxyphosphinyl] amino] pentanedioic acid;
- 2- [[(2-naphthyl) butylhydroxyphosphinyl] amino] pentanedioic acid;
- 25 2- [[(phenylprop-2-enyl) hydroxyphosphinyl] amino] pentane-dioic acid;

SUBSTITUTE SHEET (RULE 26)

264

- 2 - [[benzylhydroxyphosphinyl]amino]pentanedioic acid;
2 - [[[2-fluorobenzyl)hydroxyphosphinyl]amino] -2-pentane-
dioic acid;
2 - [[[(hydroxy)phenylmethyl)hydroxyphosphinyl]amino] -
5 pentanedioic acid;
2 - [[[3-methylbenzyl)hydroxyphosphinyl]amino]pentanedioic
acid;
2 - [[[4-fluorophenyl)hydroxyphosphinyl]amino]pentanedioic
acid;
10 2 - [(phosphono)amino]pentanedioic acid;
2 - [[[3-trifluoromethylbenzyl)hydroxyphosphinyl]amino] -
pentanedioic acid;
2 - [(methylhydroxyphosphinyl)amino]hexanedioic acid;
2 - [(benzylhydroxyphosphinyl)amino]hexanedioic acid;
15 2 - [(methylhydroxyphosphinyl)amino]heptanedioic acid;
2 - [(benzylhydroxyphosphinyl)amino]heptanedioic acid;
2 - [(methylhydroxyphosphinyl)amino]octanedioic acid;
2 - [(benzylhydroxyphosphinyl)amino]octanedioic acid;
2 - [(methylhydroxyphosphinyl)amino]nonanedioic acid;
20 2 - [(benzylhydroxyphosphinyl)amino]nonanedioic acid;
2 - [(methylhydroxyphosphinyl)amino]decanedioic acid;
2 - [(benzylhydroxyphosphinyl)amino]decanedioic acid;
3 - [[[2-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;
25 3 - [[[3-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic
acid;

SUBSTITUTE SHEET (RULE 26)

265

- 3-[[(4-pyridyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[(3-pyridyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;
- 5 3-[[(3-pyridyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[(tetrahydrofuranyl)methylhydroxyphosphinyl]amino] - pentanedioic acid;
- 3-[[(tetrahydrofuranyl)ethylhydroxyphosphinyl]amino] -
10 pentanedioic acid;
- 3-[[(tetrahydrofuranyl)propylhydroxyphosphinyl]amino] - pentanedioic acid;
- 3-[[(2-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 15 3-[[(3-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[(4-indolyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[(3-indolyl)ethylhydroxyphosphinyl]amino]pentanedioic
20 acid;
- 3-[[(3-indolyl)propylhydroxyphosphinyl]amino]pentanedioic acid;
- 3-[[(2-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;
- 25 3-[[(3-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

SUBSTITUTE SHEET (RULE 26)

266

3 - [[(4-thienyl)methylhydroxyphosphinyl]amino]pentanedioic acid;

3 - [[(3-thienyl)ethylhydroxyphosphinyl]amino]pentanedioic acid;


5 3 - [[(3-thienyl)propylhydroxyphosphinyl]amino]pentanedioic acid; and

pharmaceutically acceptable salts and hydrates thereof.

SUBSTITUTE SHEET (RULE 26)

1 / 1 4


FIG.1
COMPOUND 3 IS NEUROPROTECTIVE IN A
CELL CULTURE MODEL OF STROKE



SUBSTITUTE SHEET (RULE 26)

2 / 1 4


FIG.2
NAAG TOXICITY



SUBSTITUTE SHEET (RULE 26)

3 / 1 4


FIG.3

NAAG TOXICITY BLOCKED
BY COMPOUND 3

SUBSTITUTE SHEET (RULE 26)


4 / 1 4

FIG.4
COMPOUND 3 IS NEUROPROTECTIVE WHEN
ADMINISTERED POST-ISCHEMIA



SUBSTITUTE SHEET (RULE 26)

5 / 1 4


FIG.5**COMPOUND 3 IS NEUROPROTECTIVE
IN RAT MCAO MODEL OF STROKE**

SUBSTITUTE SHEET (RULE 26)

6 / 1 4

FIG. 6

COMPOUND 3 IS NEUROPROTECTIVE IN RAT MCAO
MODEL OF STROKE WHEN ADMINISTERED UP
TO 2 HOURS POST - OCCLUSION




SUBSTITUTE SHEET (RULE 26)

7 / 1 4


FIG.7

STRIATUM




SUBSTITUTE SHEET (RULE 26)

FIG.8
PARIETAL CORTEX



9 / 1 4

FIG. 9
FRONTAL CORTEX



SUBSTITUTE SHEET (RULE 26)

10 / 14

GPI 5000 Administration Following Sciatic Nerve Crush



Vehicle Polymer

FIG. 10A



GPI 5000 Polymer
2 ug drug / day


FIG. 10B

SUBSTITUTE SHEET (RULE 26)

11 / 14

FIG. 11

COMPOUND 3 PROTECTS AGAINST MPTP - TOXICITY IN CD1 MICE




SUBSTITUTE SHEET (RULE 26)

12 / 14


FIG.12

COMPOUND 3 IS NEUROPROTECTIVE IN A
DYNORPHIN - INDUCED SPINAL CORD INJURY MODEL



SUBSTITUTE SHEET (RULE 26)


13 / 14

FIG.13**COMPOUND 3 IS NEUROPROTECTIVE IN A
SPINAL CORD CULTURE MODEL OF ALS**

SUBSTITUTE SHEET (RULE 26)

14 / 14

FIG.14
COMPOUND 3 IS NEUROPROTECTIVE IN SPINAL CORD ORGANOTYPIC CULTURE MODEL OF ALS:
DOSE RESPONSE STUDY



SUBSTITUTE SHEET (RULE 26)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/14344

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 31/66; 31/41

US CL : 514/114, 131, 364, 376, 747, 659, 921

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/114, 131, 364, 376, 747, 659, 921

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
CAS ONLINE (STN EXPRESS)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,143,908 A (PARSONS et al.) 01 September 1992, column 2, line 20.	1-34
X	US 4,885,136 A (KATAYAMA et al.) 05 December 1989, column 4, lines 10, 20 and 24.	1-34
X	US 5,500,420 A (MAIESE et al.) 19 March 1996, claims 1, 10, and 15.	1-4; 35-42
Y	VORNOV, James J., Toxic NMDA-Receptor Activation Occurs During Recovery in a Tissue Culture Model of Ischemia, Journal of Neurochemistry, Vol. 65, No. 4, October 1993, pgs 1681-1691, especially pages 1686-1689.	1-108



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

23 SEPTEMBER 1997

Date of mailing of the international search report

23 JAN 1998

Name and mailing address of the ISA/IJS
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

KEITH MACMILLAN

Telephone No. (703) 308-1235

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US97/14344

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SLUSHER, Barbara Stauch, NAALADase: A Potential Regulator of Synaptic Glutamate, ZENECA Pharmaceuticals Group, Department of Psychology, Wilmington, Delaware, Spring 1994, pages 2-4, especially page 2.	1-108

Form PCT/ISA/210 (continuation of second sheet)(July 1992)*